

The structure of the category of parabolic equations. I

Marina Prokhorova
pmf@imm.uran.ru

Krasovskii Institute of Mathematics and Mechanics (Yekaterinburg, Russia)
Ural Federal University (Yekaterinburg, Russia)

Abstract

This is the first part of a series consisting of two papers. In this paper we define the category of partial differential equations. Special cases of morphisms from an object (equation) are symmetries of the equation and reductions of the equation by a symmetry groups, but there are many other morphisms. We develop a special-purpose language for description and study of the internal structure of this category. We are mostly interested in a subcategory that arises from second order parabolic equations on arbitrary manifolds. In the second part of the series we study the internal structure of this subcategory in detail.

Introduction

This series of two papers is a revised version of the preprint [8]. Here we continue an investigation started in [6] and [7], where the category \mathcal{PDE} of partial differential equations was defined and its study was began.

For easier introduction, let us define first the category \mathcal{PDE}_0 , which is a full subcategory of \mathcal{PDE} .

Let $\pi: N \rightarrow M$ be a smooth fiber bundle, E be a subset of k -jet bundle $J^k(\pi)$. E could be considered as a k -th order partial differential equation for sections of π ; namely, $s \in \Gamma\pi$ is a solution of E if k -th prolongation $j^k(s) \in \Gamma(J^k(\pi) \rightarrow M)$ is contained in E . Here $\Gamma\pi$ denotes the space of smooth sections of π .

Let $\pi: N \rightarrow M, \pi': N' \rightarrow M'$ be smooth fiber bundles. Suppose $F: \pi \rightarrow \pi'$ is a smooth bundle morphism with the additional property that F is a diffeomorphism on the fibers (see Fig. 1). Then F induces the map $F^*: \Gamma\pi' \rightarrow \Gamma\pi$; denote by $\Gamma_F\pi$ its image. We say that a section of π is F -projectable if it is contained in $\Gamma_F\pi$. If F is surjective, then F^* is injective, so it defines the map from $\Gamma_F\pi$ to $\Gamma\pi'$. If additionally F is submersive, then it defines the map $F^k: J^k_F(\pi) \rightarrow J^k(\pi')$, where $J^k_F(\pi) = F^*J^k(\pi')$ is the bundle of k -jets of F -projectable sections of π (see Fig. 2).

Now we are ready to define \mathcal{PDE}_0 . Its objects are pairs $(\pi: N \rightarrow M, E \subset J^k(\pi))$, $k \in \mathbb{N}$, and morphisms from an object $(\pi: N \rightarrow M, E \subset J^k(\pi))$ to an object $(\pi': N' \rightarrow M', E' \subset J^k(\pi'))$ are smooth bundle morphisms $F: \pi \rightarrow \pi'$ satisfying the following conditions:

1. F defines surjective submersion $M \rightarrow M'$.
2. The diagram Fig. 1 is a pullback square in the category of smooth manifolds, that is the map $F: \pi^{-1}(x) \rightarrow \pi'^{-1}(Fx)$ is a diffeomorphism for any $x \in M$.
3. $E \cap J^k_F(\pi) = (F^k)^{-1}(E')$.




Figure 1:




Figure 2:

Copyright © by the paper's authors. Copying permitted for private and academic purposes.

In: A.A. Makhnev, S.F. Pravdin (eds.): Proceedings of the 47th International Youth School-conference "Modern Problems in Mathematics and its Applications", Yekaterinburg, Russia, 02-Feb-2016, published at <http://ceur-ws.org>

If $F: (\pi, E) \rightarrow (\pi', E')$ is a morphism in \mathcal{PDE}_0 , then F^* defines a bijection between the set of all solutions of E' and the set of all F -projectable solutions of E .

In Section 2 we define a bigger category \mathcal{PDE} , whose objects are pairs (N, E) with N a smooth manifold and E a subset of the bundle $J_m^k(N)$ of k -jets of m -dimensional submanifolds of N , and whose morphisms from (N, E) to (N', E') are maps $N \rightarrow N'$ satisfying some analogue of conditions (1-3) above. By definition, the solutions of an equation E are smooth m -dimensional non-vertical integral manifolds of the Cartan distribution on $J_m^k(N)$, which are contained in E . Particularly, the set of solutions includes all m -dimensional submanifolds $L \subset N$ such that the k -th prolongation $j^k(L) \subset J_m^k(N)$ is contained in E . Any morphism $F: (N, E) \rightarrow (N', E')$ of \mathcal{PDE} defines a bijection between the set of all solutions of E' and the set of all F -projectable solutions of E in the same manner as for \mathcal{PDE}_0 .

The category \mathcal{PDE} generalizes the notion of symmetry group in two directions:

1. Automorphisms group of an object (N, E) in \mathcal{PDE} is the symmetry group of the equation E .
2. For a symmetry group G of E the natural projection $N \rightarrow N/G$ defines the morphism $(N, E) \rightarrow (N/G, E/G)$ in \mathcal{PDE} . Here E/G is the equation describing G -invariant solutions of E .

Note that morphisms of \mathcal{PDE} go beyond morphisms of this kind.

In Sections 2 and 6 we discuss the relations between our approach to the factorization of PDE and the other approaches.

Then we discuss the possibility of the introduction of a certain structure in \mathcal{PDE} formed by a lattice of subcategories. These subcategories may be obtained by restricting to equations of specific kind (for example, elliptic, parabolic, hyperbolic, linear, quasilinear equations etc.) or to the morphisms of specific kind (for example, morphisms respecting the projection of N on a base manifold M as in \mathcal{PDE}_0) or both. When we interested in solutions of some equation it is useful to look for its quotient objects because every quotient object gives us a class of solutions of the original equation. It may happen that the position of an object in the lattice gives information on the morphisms from the object and/or on the kind of the simplest representatives of quotient objects. In Section 4 we develop a special-purpose language for description and study of such situations. We introduce a number of partial orders on the class of all subcategories of fixed category and depict these orders by various arrows (see Table 1 and Fig. 3). For instance, we say that a subcategory \mathcal{C}_1 is closed in a category \mathcal{C} and depict $\mathcal{C} \bullet \longrightarrow \mathcal{C}_1$ if every morphism in \mathcal{C} with source from \mathcal{C}_1 is a morphism in \mathcal{C}_1 ; we say that \mathcal{C}_1 is plentiful in \mathcal{C} and depict $\mathcal{C} - - \triangleright \mathcal{C}_1$ if for every $\mathbf{A} \in \text{Ob}_{\mathcal{C}_1}$ and for every quotient object of \mathbf{A} in \mathcal{C} there exists a representative of this quotient object in \mathcal{C}_1 ; and so on.

We use this language in the second part of this series [9] for a detail study of the full subcategory \mathcal{PE} of \mathcal{PDE} that arises from second order parabolic equations posed on arbitrary manifolds, but we hope that our approach based on category theory may be useful for other types of PDE as well. An object of \mathcal{PE} is an equation for an unknown function $u(t, x)$, $x \in X$ having the form $u_t = \sum_{i,j} b^{ij}(t, x, u)u_{ij} + \sum_{i,j} c^{ij}(t, x, u)u_i u_j + \sum_i b^i(t, x, u)u_i + q(t, x, u)$ in local coordinates (x^i) on X , where X is a smooth manifold. We prove that every morphism in \mathcal{PE} is of the form $(t, x, u) \mapsto (t'(t), x'(t, x), u'(t, x, u))$ (Theorem 1). Particularly, \mathcal{PE} appears to be a subcategory of \mathcal{PDE}_0 .

The use of the structure of \mathcal{PE} developed in the second paper [9] is illustrated there on the example of the reaction-diffusion equation

$$u_t = a(u) (\Delta u + \eta \nabla u) + q(x, u), \quad x \in X, \quad t \in \mathbb{R}, \quad (1)$$

posed on a Riemannian manifold X equipped with a vector field η . There are two exceptional cases: $a(u) = e^{\lambda u} H(u)$ and $a(u) = (u - u_0)^\lambda H(\ln(u - u_0))$, where $H(\cdot)$ is a periodic function; in these cases there are more morphisms than in a regular case. If only function $a(u)$ does not belong to one of these two exceptional classes, then every morphism from equation (1) may be transformed by an isomorphism (i.e. by a global change of variables) of the quotient equation to a ‘‘canonical’’ morphism of very simple kind so that the ‘‘canonical’’ quotient equation has the same form as (1) with the same function $a(u)$ but is posed on another Riemannian manifold X' , $\dim X' \leq \dim X$.

1 The ‘‘small’’ category \mathcal{PDE}_0 of partial differential equations

Let M, K be smooth manifolds. A system E of k -th order partial differential equations for a function $u: M \rightarrow K$ is given as a system of equations $\Phi^l(x, u, \dots, u^{(k)}) = 0$ involving x, u and the derivatives of u with respect to x up to order k , where $x = (x^1, \dots, x^m)$ are local coordinates on M and $u = (u^1, \dots, u^j)$ are local coordinates

on K . Further we will use the words “partial differential equation”, “PDE” or “equation” instead of “a partial differential equation or a system of partial differential equations” for short.

Recall some things about jets and related notions. The k -jet of a smooth function $u: M \rightarrow K$ at a point $x \in M$ is the equivalence class of smooth functions $M \rightarrow K$ whose value and partial derivatives up to k -th order at x coincide with the ones of u . All k -jets of all smooth functions $M \rightarrow K$ form the smooth manifold $J^k(M, K)$, and the natural projection $\pi^k: J^k(M, K) \rightarrow J^0(M, K) = M \times K$ defines a smooth vector bundle over $M \times K$, which is called k -jet bundle. For every function $u: M \rightarrow K$, its k -th prolongation $j^k(u): M \rightarrow J^k(M, K)$ is naturally defined. k -th order PDE for functions acting from M to K can be considered as a subset E of $J^k(M, K)$; solutions of E are functions $u: M \rightarrow K$ such that the image of $j^k(u)$ is contained in E .

In more general situation we have a smooth fiber bundle $\pi: N \rightarrow M$ instead of a projection $M \times K \rightarrow M$, and sections $s: M \rightarrow N$ instead of functions $u: M \rightarrow K$. Denote by $\Gamma\pi$ the space of smooth sections of π ; recall that a section of π is a map $s: M \rightarrow N$ such that $\pi \circ s$ is the identity. Definitions of the k -jet bundle $\pi^k: J^k(\pi) \rightarrow J^0(\pi) = N$ and of the k -th prolongation $j^k: \Gamma\pi \rightarrow \Gamma(\pi \circ \pi^k: J^k(\pi) \rightarrow M)$ are the same as those for functions. Let E be a subset of $J^k(\pi)$; then E can be considered as a k -th order partial differential equation for sections of π , that is $s \in \Gamma\pi$ is a solution of E if the image of $j^k(s)$ is contained in E .

Let $\pi: N \rightarrow M$, $\pi': N' \rightarrow M'$ be smooth fiber bundles. Let $F: \pi \rightarrow \pi'$ be a smooth bundle morphism with the additional property that F is a diffeomorphism on the fibers (see Fig. 1). F induces the map $F^*: \Gamma\pi' \rightarrow \Gamma\pi$; denote by $\Gamma_F\pi$ its image. We say that a section of π is F -projectable if it is contained in $\Gamma_F\pi$. If F is surjective, then F^* is injective, so it defines the map $F_\#: \Gamma_F\pi \rightarrow \Gamma\pi'$. If additionally F is submersive, then it defines the map $F^k: J_F^k(\pi) \rightarrow J^k(\pi')$, where $J_F^k(\pi) = F^*J^k(\pi')$ is the bundle of k -jets of F -projectable sections of π (see Fig. 2). Recall that a map F is called a submersion if $dF: T_xN \rightarrow T_{F(x)}N'$ is surjective at each point $x \in N$.

Definition 1. Let $\pi: N \rightarrow M$, $\pi': N' \rightarrow M'$ be smooth fiber bundles, E be a subset of $J^k(\pi)$, $F: \pi \rightarrow \pi'$ be a smooth bundle morphism. We say that F is admitted by E if the following conditions are satisfied:

1. F is a surjective submersion.
2. The diagram Fig. 1 is a pullback square in the category of smooth manifolds, that is the map $F: \pi^{-1}(x) \rightarrow \pi'^{-1}(Fx)$ is a diffeomorphism for any $x \in M$.
3. $E \cap J_F^k(\pi) = (F^k)^{-1}(E')$ fore some subset E' of $J^k(\pi')$.

If this is the case, then we say that E' is F -projection of E .

It turns out that the language of category theory is very convenient for our study of PDE. Recall that a category \mathcal{C} consists of a collection of *objects* $\text{Ob}_{\mathcal{C}}$, a collection of *morphisms* (or *arrows*) $\text{Hom}_{\mathcal{C}}$ and four operations. The first two operations associate with each morphism F of \mathcal{C} its *source* and its *target*, both of which are objects of \mathcal{C} . The remaining two operations are an operation that associates with each object \mathbf{C} of \mathcal{C} an *identity morphism* $\text{id}_{\mathbf{C}} \in \text{Hom}_{\mathcal{C}}$ and an operation of *composition* that associates to any pair (F, G) of morphisms of \mathcal{C} such that the source of F coincides with the target of G another morphism $F \circ G$, their composite. These operations should satisfy some natural axioms [3, sec. I.1].

Definition 2. \mathcal{PDE}_0 is the category whose objects are pairs $(\pi: N \rightarrow M, E \subset J^k(\pi))$ with π being a smooth viber bundle, $k \in \mathbb{N}$, and morphisms from an object $(\pi: N \rightarrow M, E \subset J^k(\pi))$ to an object $(\pi': N' \rightarrow M', E' \subset J^k(\pi'))$ are smooth bundle morphisms $F: \pi \rightarrow \pi'$ admitted by E such that E' is the F -projection of E .

If $F: (\pi, E) \rightarrow (\pi', E')$ is a morphism in \mathcal{PDE}_0 , then F^* defines a bijection between the set of all solutions of E' and the set of all F -projectable solutions of E .

2 The category \mathcal{PDE} of partial differential equations

In this section we define the category \mathcal{PDE} of partial differential equations, whose objects are pairs (N, E) such that N is a smooth manifold and E is a subset of the bundle $J_m^k(N)$ of k -jets of m -dimensional submanifolds of N .

Let N be a C^r -smooth manifold, $0 < m < \dim N$. The jet bundle $\pi^k: J_m^k(N) \rightarrow N$ is a fiber bundle with the fiber $J_m^k(N)|_x$ over $x \in N$, where $J_m^k(N)|_x$ is the set of equivalence classes of smooth m -dimensional submanifolds L of N passing through x under the equivalence relation of k -th order contact in x .

The k -jet of a k -smooth m -dimensional submanifold L over $x \in L$ is the equivalence class from $J_m^k(N)|_x$ determined by L . Thus we have the prolongation map $j^k: L \rightarrow J_m^k(N)$ taking each point $x \in L$ to the k -jet

of L over x (so it is the section of the fiber bundle $J_m^k(N)$ restricted to $L \subset N$). For every $k > l \geq 0$ the natural projection $\pi^{k,l}: J_m^k(N) \rightarrow J_m^l(N)$ maps the k -jet of L to the l -jet of L over x for every m -dimensional submanifold L of N and every $x \in L$.

For a submanifold L of N the differential of the prolongation map $j^k: L \rightarrow J_m^k(N)$ takes the tangent bundle TL to the tangent bundle $TJ_m^k(N)$. The closure of the union of the images of TL in $TJ_m^k(N)$ when L runs over all m -dimensional submanifolds of N is the vector subbundle of $TJ_m^k(N)$; it is called the Cartan distribution on $J_m^k(N)$.

Let E be a submanifold of $J^k(\pi)$, $\pi: N \rightarrow M$, $m = \dim M$. The graph of a section is an m -dimensional submanifold of N , so $J^k(\pi)$ is an open subspace of $J_m^k(N)$ and E could be considered as a submanifold of $J_m^k(N)$. The extended version of E is defined as the closure of E in $J_m^k(N)$ [4, p. 222]. Since we don't plan to consider infinitesimal properties of E in contrast to the Lie group analysis of PDE, we would consider any subset E of $J_m^k(N)$ as a partial differential equations. By definition, *solutions* of such an equation are smooth m -dimensional non-vertical integral manifolds of the Cartan distribution on $J_m^k(N)$ that are contained in E . Note that for any m -dimensional submanifold L of N its prolongation $j^k(L)$ is a non-vertical integral manifold of the Cartan distribution on $J_m^k(N)$. Therefore, if $E \subset J_m^k(N)$ is obtained from a traditional PDE as it was described above, and if L is the graph of a section u of π , then L is a solution of E in the above sense if and only if u is a solution of the corresponding traditional PDE in the traditional sense. In addition there is allowed the possibility of both multi-valued solutions and solutions with infinite derivatives (see [4, sec. 3.5] for the details). Wherever we write concrete equation in the traditional form below we mean the extended version of this equations, that is the closure of the corresponding set in $J_m^k(N)$.

Now let us introduce some auxiliary notation.

Let $F: N \rightarrow N'$ be a map. We say that $L \subset N$ is F -projected if $L = F^{-1}(F(L))$. Note that if F is a surjective submersion and L is an F -projectable submanifold of N , then $L' = F(L)$ is a submanifold of N' .

Let N, N' be C^r -smooth manifolds, $0 < m < \dim N$. Let $F: N \rightarrow N'$ be a surjective submersion of smoothness class C^s , $k \leq s \leq r$.

Definition 3. F -projectable jet bundle $J_{m,F}^k(N)$ is the submanifold of $J_m^k(N)$ consisting of k -jets of all m -dimensional F -projectable submanifolds of N .

We write $J_F^k(N)$ instead of $J_{m,F}^k(N)$ if the value of m is clear from context.

There is natural isomorphism between the bundles $J_{m,F}^k(N)$ and $F^*J_{m'}^k(N')$ over N , where $F^*J_{m'}^k(N') = J_{m'}^k(N') \times_{N'} N$ is the pullback of $J_{m'}^k(N')$ by F , $\dim N - m = \dim N' - m'$. Therefore we can lift F to the map $F^k: J_{m,F}^k(N) \rightarrow J_{m'}^k(N')$ by the following natural way:

1) Suppose $\vartheta \in J_{m,F}^k(N)$. Take an arbitrary F -projectable submanifold L of N such that the k -th prolongation of L pass through ϑ (that is the k -jet of L over $\pi^k(\vartheta)$) is ϑ .

2) Assign to ϑ the point $\vartheta' \in J_{m'}^k(N')$, where ϑ' is the k -jet of the submanifold $L' = F(L) \subset N'$ over $F \circ \pi^k(\vartheta)$.

Definition 4. Let $E \subset J_m^k(N)$. Let $F: N \rightarrow N'$ be a smooth surjective submersion. We say that F is *admitted by* E if the intersection $E \cap J_{m,F}^k(N)$ is F^k -projectable subset of $J_{m,F}^k(N)$ (see the left diagram on Fig. 3). Equivalently, $E \cap J_{m,F}^k(N)$ is the inverse image $(F^k)^{-1}(E')$ of some $E' \subset J_{m'}^k(N')$; we say that E' is F -projection of E .




Figure 3: Morphisms of \mathcal{PDE} (left diagram) and morphisms of $\mathcal{PDE}_{\text{ext}}$ (right diagram).

Definition 5. The *category of partial differential equations* \mathcal{PDE} is defined as follows:

- objects of \mathcal{PDE} are pairs (N, E) , where N is a smooth manifold, E is a subset of $J_m^k(N)$ for some integer $k, m \geq 1$;
- morphisms of \mathcal{PDE} with a source $\mathbf{A} = (N, E)$ are all surjective submersions $F: N \rightarrow N'$ admitted by E ; target of such morphism is (N', E') where E' is F -projection of E ;
- the identity morphism from \mathbf{A} is the identity mapping of N , and the composition of morphisms is the composition of appropriate maps.

If $F: (N, E) \rightarrow (N', E')$ is a morphism in \mathcal{PDE} , then F^* defines a bijection between the set of all solutions of E' and the set of all F^k -projectable solutions of E .

In [5] the author defined the following notion of a map admitted by a pair of equations: a map $F: N \rightarrow N'$ is admitted by an ordered pair of equations $(\mathbf{A}, \mathbf{A}')$, $\mathbf{A} = (N, E)$, $\mathbf{A}' = (N', E')$ if for any $L' \subset N'$ the following two conditions are equivalent:

- L' is the graph of a solution of E' ,
- $F^{-1}(L')$ is the graph of a solution of E .

However, we are not happy with this definition; in particular, because it deals only with global solutions of E . Therefore, here we define the notion of map admitted by an equation in terms of (locally defined) jet bundles.

Remark 1. Let $\mathbf{A} = (N, E)$ be an object of \mathcal{PDE} . Then its automorphism group $\text{Aut}(\mathbf{A})$ is the symmetry group for the equation E .

Remark 2. Suppose G is a subgroup of the symmetry group of E such that N/G is a smooth manifold. Then the natural projection $N \rightarrow N/G$ defines the morphism $(N, E) \rightarrow (N/G, E/G)$ in \mathcal{PDE} . Here E/G is the equation describing G -invariant solutions of E .

Therefore, reduction of E by subgroups of $\text{Aut}(\mathbf{A})$ defines a part of nontrivial morphisms from \mathbf{A} . But the class of all morphisms from \mathbf{A} is significantly richer than the class of morphisms arising from reduction by subgroups of $\text{Aut}(\mathbf{A})$. Let $\text{Sol}(\mathbf{A})$ be the set of all solutions of \mathbf{A} , that is of all smooth m -dimensional non-vertical integral manifolds of the Cartan distribution on $J_m^k(N)$ that are contained in E . In general, the subset $F^*(\text{Sol}(\mathbf{A}')) = \{F^{-1}(L') : L' \in \text{Sol}(\mathbf{A}')\} \subseteq \text{Sol}(\mathbf{A})$ of solutions of \mathbf{A} arising from a morphism $F: \mathbf{A} \rightarrow \mathbf{A}'$ can *not* be represented as a set of solutions that are invariant under some subgroup of $\text{Aut}(\mathbf{A})$. In particular, $F^*(\text{Sol}(\mathbf{A}'))$ can be the set of G -invariant solutions, where G is a transformation group that is not necessarily a symmetry group of E . Moreover, for a morphism $F: \mathbf{A} \rightarrow \mathbf{A}'$ it may occur that for every nontrivial diffeomorphism g of N there is an element in $F^*(\text{Sol}(\mathbf{A}'))$ that is not g -invariant. More detailed discussion is given in Section 6; see also [6, sections 9, 10], [7, sections 7–9].

Our approach is conceptually close to the approach developed in [1] that deals with control systems. If we set aside the control part and look at this approach relative to ordinary differential equations, then we get the category of ordinary differential equations, whose objects are ODE systems of the form $\dot{x} = \xi$, $x \in X$, where X is a manifold equipped with a vector field ξ , and morphism from a system \mathbf{A} to a system \mathbf{A}' is a smooth map F from the phase space X of \mathbf{A} to the phase space X' of \mathbf{A}' that projected ξ to ξ' . In other words, F is a morphism if it transforms solutions (phase trajectories) of \mathbf{A} to solutions of \mathbf{A}' : $F_*(\text{Sol}(\mathbf{A})) = \text{Sol}(\mathbf{A}')$.

By contrast, we deal with pullbacks of the solutions of the quotient equation \mathbf{A}' to the solutions of the original equation \mathbf{A} . In our approach, the number of dependent variables in the reduced PDE remains the same, while the number of independent variables is not increased. Thus, in the approach proposed, the quotient object notion is an analogue of the sub-object notion (in terminology of [1, sec. 5.1]) with respect to the information about the solutions of the given equation; however, it is similar to the quotient object notion with respect to interrelations between the given and reduced equations.

Note also that the above described category of ODE from [1] is isomorphic to a certain subcategory of \mathcal{PDE} . Namely, consider the following subcategory \mathcal{PDE}_1 of \mathcal{PDE} :

- objects of \mathcal{PDE}_1 are pairs (N, E) , where $N = X \times \mathbb{R}$, E is a first order linear PDE of the form $L_\xi u = 1$ for unknown function $u: X \rightarrow \mathbb{R}$, $\xi \in TX$;
- morphisms of \mathcal{PDE}_1 are morphisms of \mathcal{PDE} of the form $(x, u) \mapsto (x'(x), u)$.

One can easily see that the category of ODE from [1] is isomorphic to \mathcal{PDE}_1 : the object $L_\xi u = 1$ corresponds to the object $\dot{x} = \xi$, and the morphism $(x, u) \mapsto (x'(x), u)$ corresponds to the morphism $x \mapsto x'(x)$.

- the identity morphism from \mathbf{A} is $(\Delta = J_m^k(N), \tilde{F} = \text{id}_N)$;
- composition of $(\Delta \subset J_m^k(N), \tilde{F}: \Delta \rightarrow J_m^{k'}(N'))$ and $(\Delta' \subset J_m^{k''}(N''), \tilde{F}': \Delta' \rightarrow J_m^{k''}(N''))$ is $(\tilde{F}^{-1}(\Delta'), \tilde{F}' \circ \tilde{F})$.

For each integral manifold of the Cartan distribution on E' , its inverse image is an integral manifold of the Cartan distribution on E . Therefore, for each solution of E' , its pullback is a solution of E .

\mathcal{PDE} is embedded to $\mathcal{PDE}_{\text{ext}}$ by the following natural way: to the morphisms $F: N \rightarrow N'$ of \mathcal{PDE} from the equation of k -th order we assign the morphisms (Δ, \tilde{F}) of $\mathcal{PDE}_{\text{ext}}$ such that $\Delta = J_{m,F}^k(N)$, $\tilde{F} = F^k$.

4 Usage of subcategories

We start with a review of some basic definitions of category theory [3, sec. II.6]. Given a category \mathcal{C} and an object \mathbf{A} of \mathcal{C} , one may construct the category $(\mathbf{A} \downarrow \mathcal{C})$ of objects under \mathbf{A} (this is the particular case of the comma category): objects of $(\mathbf{A} \downarrow \mathcal{C})$ are morphisms of \mathcal{C} with source \mathbf{A} , and morphisms of $(\mathbf{A} \downarrow \mathcal{C})$ from one such object $F: \mathbf{A} \rightarrow \mathbf{B}$ to another $F': \mathbf{A} \rightarrow \mathbf{B}'$ are morphisms $G: \mathbf{B} \rightarrow \mathbf{B}'$ of \mathcal{C} such that $F' = G \circ F$.

Suppose \mathcal{C} is a subcategory of \mathcal{PDE} , \mathbf{A} is an object of \mathcal{C} . Then the category $(\mathbf{A} \downarrow \mathcal{C})$ of objects under \mathbf{A} describes collection of quotient equations for \mathbf{A} and their interconnection in the framework of \mathcal{C} .

To each morphism $F: \mathbf{A} \rightarrow \mathbf{B}$ with source \mathbf{A} (that is to each object of the comma category $(\mathbf{A} \downarrow \mathcal{C})$) assign the set $F^*(\text{Sol}(\mathbf{B})) \subseteq \text{Sol}(\mathbf{A})$ of such solutions of \mathbf{A} that “projected” onto underlying space of \mathbf{B} (space of dependent and independent variables). We can identify such morphisms that generated the same sets of solutions of \mathbf{A} , that is identify isomorphic objects of the comma category $(\mathbf{A} \downarrow \mathcal{C})$.

Describe the situation more explicitly. An equivalence class of epimorphisms with source \mathbf{A} is called a quotient object of \mathbf{A} , where two epimorphisms $F: \mathbf{A} \rightarrow \mathbf{B}$ and $F': \mathbf{A} \rightarrow \mathbf{B}'$ are equivalent if $F' = I \circ F$ for some isomorphism $I: \mathbf{B} \rightarrow \mathbf{B}'$ [3, sec. V.7]. If $F: \mathbf{A} \rightarrow \mathbf{B}$ and $F': \mathbf{A} \rightarrow \mathbf{B}'$ are equivalent, then they lead to the same subsets of the solutions of \mathbf{A} : $F^*(\text{Sol}(\mathbf{B})) = F'^*(\text{Sol}(\mathbf{B}'))$. So if we interested only in the sets of the solutions of \mathbf{A} , then all representatives of the same quotient object have the same rights.

Therefore, the following problems naturally arise:

- to study all morphisms with given source,
- to choose a “simplest” representative from every equivalence class, or to choose representative with the simplest target (that is the simplest quotient equation).

In order to deal with these problems we develop in this paper a special-purpose language.

Let us introduce a number of partial orders on the class of all categories to describe arising situations. First of all, we define a few types of subcategories.

Definition 7. Suppose \mathcal{C} is a category, \mathcal{C}_1 is a subcategory of \mathcal{C} .

- \mathcal{C}_1 is called a *wide* subcategory of \mathcal{C} if all objects of \mathcal{C} are objects of \mathcal{C}_1 .
- \mathcal{C}_1 is called a *full* subcategory of \mathcal{C} if every morphism in \mathcal{C} with source and target from \mathcal{C}_1 is a morphism in \mathcal{C}_1 .
- We say that \mathcal{C}_1 is *full under isomorphisms* in \mathcal{C} if every isomorphism in \mathcal{C} with source and target from \mathcal{C}_1 is an isomorphism in \mathcal{C}_1 .
- We say that \mathcal{C}_1 is *closed* in \mathcal{C} if every morphism in \mathcal{C} with source from \mathcal{C}_1 is a morphism in \mathcal{C}_1 . (Note that every subcategory that is closed in \mathcal{C} is full in \mathcal{C} .)
- We say that \mathcal{C}_1 is *closed under isomorphisms* in \mathcal{C} if every isomorphism in \mathcal{C} with source from \mathcal{C}_1 is an isomorphism in \mathcal{C}_1 .
- We say that \mathcal{C}_1 is *dense* in \mathcal{C} if every object of \mathcal{C} is isomorphic in \mathcal{C} to an object of \mathcal{C}_1 .
- We say that \mathcal{C}_1 is *plentiful* in \mathcal{C} if for every morphism $F: \mathbf{A} \rightarrow \mathbf{B}$ in \mathcal{C} , $\mathbf{A} \in \text{Ob}_{\mathcal{C}_1}$, there exists an isomorphism $I: \mathbf{B} \rightarrow \mathbf{C}$ in \mathcal{C} such that $I \circ F \in \text{Hom}_{\mathcal{C}_1}$ (in other words, for every quotient object of \mathbf{A} in \mathcal{C} there exists a representative of this quotient object in \mathcal{C}_1). Such morphism $I \circ F$ we call \mathcal{C}_1 -**canonical** for F .
- We say that \mathcal{C}_1 is *fully dense* (*fully plentiful*) in \mathcal{C} if \mathcal{C}_1 is a full subcategory of \mathcal{C} and \mathcal{C}_1 is dense (plentiful) in \mathcal{C} .

The first two parts of this definition are standard notions of category theory, while the notions of the other parts are introduced here for the sake of description of the structure of \mathcal{PDE} .

Remark 3. Using the notion of “the category of objects under \mathbf{A} ”, we can define the notions of closed subcategory and plentiful subcategory by the following way:

- \mathcal{C}_1 is closed in \mathcal{C} if for each $\mathbf{A} \in \text{Ob}_{\mathcal{C}_1}$ the category $(\mathbf{A} \downarrow \mathcal{C}_1)$ is wide in $(\mathbf{A} \downarrow \mathcal{C})$.
- \mathcal{C}_1 is plentiful in \mathcal{C} if for each $\mathbf{A} \in \text{Ob}_{\mathcal{C}_1}$ the category $(\mathbf{A} \downarrow \mathcal{C}_1)$ is dense in $(\mathbf{A} \downarrow \mathcal{C})$.

Remark 4. \mathcal{C}_1 is fully dense in \mathcal{C} if and only if the embedding functor $\mathcal{C}_1 \rightarrow \mathcal{C}$ defines an equivalence of these categories.

Choose some category \mathcal{U} , which is big enough to contain all needful for us categories as it's subcategories. For our purposes $\mathcal{U} = \mathcal{PDE}$ is sufficient.

Define the category \mathcal{U}_{\geq} , whose objects are subcategories \mathcal{C} of \mathcal{U} , and a collection $\text{Hom}_{\mathcal{U}_{\geq}}(\mathcal{C}_1, \mathcal{C}_2)$ of morphisms from \mathcal{C}_1 to \mathcal{C}_2 is a one-element set if \mathcal{C}_2 is subcategory of \mathcal{C}_1 and empty otherwise, so an arrow from \mathcal{C}_1 to \mathcal{C}_2 in \mathcal{U}_{\geq} means that \mathcal{C}_2 is the subcategory of \mathcal{C}_1 . Let $\mathcal{U}_=$ be the discrete wide subcategory of \mathcal{U}_{\geq} , that is objects of $\mathcal{U}_=$ are all subcategories \mathcal{C} of \mathcal{U} , and the only morphisms are identities, so \mathcal{C}_1 and \mathcal{C}_2 are connected by arrow in $\mathcal{U}_=$ only if $\mathcal{C}_1 = \mathcal{C}_2$.

Definition 8. Suppose $\mathcal{C}_1, \mathcal{C}_2$ are subcategories of \mathcal{U} . A subcategory of \mathcal{U} , whose objects are objects of \mathcal{C}_1 and \mathcal{C}_2 simultaneously, and whose morphisms are morphisms of \mathcal{C}_1 and \mathcal{C}_2 simultaneously, is called an *intersection* of \mathcal{C}_1 and \mathcal{C}_2 and is denoted by $\mathcal{C}_1 \cap \mathcal{C}_2$. In other words, $\mathcal{C}_1 \cap \mathcal{C}_2$ is the fibered sum of \mathcal{C}_1 and \mathcal{C}_2 in \mathcal{U}_{\geq} .

The following proposition is obvious:

Proposition 3. Suppose \mathcal{C}_1 is closed in \mathcal{C} , and \mathcal{C}_2 is (full/closed/dense/plentiful) subcategory of \mathcal{C} ; then $\mathcal{C}_1 \cap \mathcal{C}_2$ is closed in \mathcal{C}_2 and is (full/closed/dense/plentiful) subcategory of \mathcal{C}_1 .

Now we introduce some graphic designations for various types of subcategories of \mathcal{U}_{\geq} (see Table 1). These designations will be used, particularly, for the representation of the structure of the category of parabolic equations described below.

We shall use the term “meta-category” both for the category \mathcal{U}_{\geq} and for its subcategories defined below to avoid confusion between \mathcal{U}_{\geq} and “ordinary” categories which are objects of \mathcal{U}_{\geq} ; and we shall use Gothic script for meta-categories except \mathcal{U}_{\geq} . One may view these meta-categories as a partial orders on the class of all subcategories of \mathcal{U} ; we prefer category terminology here since this allows us to use category constructions for the interrelations between various partial orders.

\Rightarrow	\mathfrak{W}	Wide
$\bullet \rightarrow$	\mathfrak{F}	Full
$\circ \rightarrow$	\mathfrak{F}_1	Full under isomorphisms
$\bullet \dashrightarrow$	\mathfrak{C}	Close
$\circ \dashrightarrow$	\mathfrak{C}_1	Close under isomorphisms
\dashrightarrow	\mathfrak{D}	Dense
\dashrightarrow	\mathfrak{P}	Plentiful

Table 1: Basic meta-categories (arrows)

Let us define wide subcategories $\mathfrak{W}, \mathfrak{F}, \mathfrak{F}_1, \mathfrak{C}, \mathfrak{C}_1, \mathfrak{D}$, and \mathfrak{P} of meta-category \mathcal{U}_{\geq} . Objects of them are categories, while arrows from \mathcal{C}_1 to \mathcal{C}_2 have a different meaning:

- in the meta-category \mathfrak{W} it means that \mathcal{C}_2 is wide subcategory of \mathcal{C}_1 ,
- in the meta-category \mathfrak{F} it means that \mathcal{C}_2 is full subcategory of \mathcal{C}_1 ,
- in the meta-category \mathfrak{F}_1 it means that \mathcal{C}_2 is full under isomorphisms in \mathcal{C}_1 ,
- in the meta-category \mathfrak{C} it means that \mathcal{C}_2 is closed subcategory of \mathcal{C}_1 ,
- in the meta-category \mathfrak{C}_1 it means that \mathcal{C}_2 is closed under isomorphisms in \mathcal{C}_1 ,
- in the meta-category \mathfrak{D} it means that \mathcal{C}_2 is dense subcategory of \mathcal{C}_1 ,

- in the meta-category \mathfrak{P} it means that \mathcal{C}_2 is plentiful subcategory of \mathcal{C}_1 ,

We shall denote the intersections of these meta-categories by the concatenations of appropriate letters, for example: $\mathfrak{F}\mathcal{D} = \mathfrak{F} \cap \mathcal{D}$. The following proposition is obvious.

Proposition 4. $\mathfrak{F}_1 \cap \mathfrak{P} = \mathfrak{F} \cap \mathfrak{P}$; $\mathcal{C}_1 \cap \mathfrak{P} = \mathcal{C}$; $\mathfrak{F} \cap \mathfrak{P} \cap \mathcal{D} = \mathfrak{F}\mathcal{D}$.

Interrelations between “basic” meta-categories \mathfrak{W} , \mathfrak{F} , \mathfrak{F}_1 , \mathcal{C} , \mathcal{C}_1 , \mathcal{D} , \mathfrak{P} and their intersections (“composed” meta-categories) are represented on Fig. 5(a). Here an arrow means the predicate “to be subcategory of”; we shall call it the “meta-arrow”. For example, meta-arrow from \mathcal{D} to \mathfrak{W} means that \mathfrak{W} is a subcategory of \mathcal{D} . In the language of “ordinary” categories this meta-arrow means that the statement “ \mathcal{C}_2 is wide in \mathcal{C}_1 ” implies that \mathcal{C}_2 is dense in \mathcal{C}_1 . Everywhere on Fig. 5(a) a pair of meta-arrows with the same target means that this meta-category (target of these meta-arrows) is the intersection of two “top” meta-categories (sources of these meta-arrows). For example, $\mathfrak{F}\mathcal{D} = \mathfrak{F}\mathfrak{P} \cap \mathfrak{P}\mathcal{D}$.

On Fig. 5(b) the same scheme is represented as on Fig. 5(a), but the letter names are replaced by the arrows of various types.




Figure 5: Interrelations between basic meta-categories (arrows) and their intersections.

Instead of investigation of all or the simplest morphisms with the given source, we want to introduce a certain structure in \mathcal{PDE} , so that the position of an object in it gives an information about the morphisms from the object and about the kind of the simplest representatives of equivalence classes of the morphisms. In the second part of this series [9] we describe such a structure for the category of parabolic equations, choosing some subcategories of \mathcal{PE} connected by the arrows from Fig. 5(b). Then we use this structure to describe the morphisms from nonlinear reaction-diffusion equation.

5 The category of parabolic equations

Let us consider the class $P(X, T, \Omega)$ of differential operators on a connected smooth manifold X , which depend additionally on a parameter t (“time”), locally having the form

$$Lu = \sum_{i,j} b^{ij}(t, x, u) u_{ij} + \sum_{i,j} c^{ij}(t, x, u) u_i u_j + \sum_i b^i(t, x, u) u_i + q(t, x, u),$$

$$x \in X, \quad t \in T, \quad u \in \Omega,$$

in some neighborhood of each point, in some (and then arbitrary) local coordinates (x^i) on X . Here subscript i denotes partial derivative with respect to x^i , quadratic form $b^{ij} = b^{ji}$ is positive definite, and $c^{ij} = c^{ji}$. Both T and Ω may be bounded, semi-bounded, or unbounded open intervals of \mathbb{R} .

Definition 9. The category \mathcal{PE} of parabolic equations of the second order is the full subcategory of \mathcal{PDE} , whose objects are pairs $\mathbf{A} = (N, E)$, $N = T \times X \times \Omega$ such that X is a connected smooth manifold, T and Ω are open

intervals, and E is an equation of the form $u_t = Lu$, $L \in P(X, T, \Omega)$ (more exactly, E is the extended version of the equation $u_t = Lu$, that is a closed submanifold of $J_{n+1}^2(T \times X \times \Omega)$, $n = \dim X$).

Example 1. Let $\Phi_k(x)$, $x \in \mathbb{R}^3 - \{0\}$ be a spherical harmonic of the k -th order. Then the map $(t, x, u) \mapsto (t, |x|, u/\Phi_k(x))$ defines the morphism in the category \mathcal{PE} from the object \mathbf{A} corresponding to equation $u_t = \Delta u$ and $X = \mathbb{R}^3 - \{0\}$, $T = \Omega = \mathbb{R}$, to the object \mathbf{A}' corresponding to equation $u'_t = u'_{x'x'} - k(k+1)x'^{-2}u'$ and $X' = \mathbb{R}_+$, $T' = \Omega' = \mathbb{R}$. One may assign to the set $\text{Sol}(\mathbf{A}')$ of all solutions of the quotient equation the set $F^*(\text{Sol}(\mathbf{A}'))$ of such solutions of the original equation that may be written in the form $u = \Phi_k(x)u'(t, |x|)$.

Example 2. The following example shows that not every endomorphism in \mathcal{PE} is an automorphism. Consider object \mathbf{A} , for which $X = S^1 = \mathbb{R} \bmod 1$, $T = \Omega = \mathbb{R}$, $E: u_t = u_{xx}$. Then the morphism from \mathbf{A} to \mathbf{A} defined by the map $(t, x, u) \mapsto (4t, 2x, u)$ has no inverse.

Theorem 1. *Every morphism in \mathcal{PE} has the form*

$$(t, x, u) \mapsto (t'(t), x'(t, x), u'(t, x, u)), \quad (2)$$

with submersive $t'(t)$, $x'(t, x)$, and $u'(t, x, u)$. Isomorphisms in \mathcal{PE} are exactly diffeomorphisms of the form (2).

Proof. Passing from the equation $u_t = Lu$ to the equation in the extended jet bundle for unknown submanifold $L \subset X \times T \times \Omega$ locally defined by the formula $f(t, x, u) = 0$, and expressing the derivatives of u by the corresponding derivatives of f , we obtain the following extended version of E :

$$f_t f_u^2 = \sum_{i,j} b^{ij} (f_{ij} f_u^2 - (f_{iu} f_j + f_{ju} f_i) f_u + f_i f_j f_{uu}) - \sum_{i,j} c^{ij} f_i f_j f_u + \sum_i b^i f_i f_u^2 - q f_u^3. \quad (3)$$

Suppose $F: \mathbf{A} \rightarrow \mathbf{A}'$ is a morphism in \mathcal{PE} , $N' = X' \times T' \times \Omega'$, and E' is defined by the equation

$$u' = \sum_{i',j'} B^{i'j'}(t', x', u') u'_{i'j'} + \sum_{i',j'} C^{i'j'}(t', x', u') u'_{i'} u'_{j'} + \sum_{i'} B^{i'}(t', x', u') u'_{i'} + Q(t', x', u').$$

Consider the extended analog of the last equation:

$$\begin{aligned} f'_{t'} f'^2_{u'} &= \sum_{i',j'} B^{i'j'} \left(f'_{i'j'} f'^2_{u'} - (f'_{i'u'} f'_{j'} + f'_{j'u'} f'_{i'}) f'_{u'} + f'_{i'} f'_{j'} f'_{u'u'} \right) \\ &\quad - \sum_{i',j'} C^{i'j'} f'_{i'} f'_{j'} f'_{u'} + \sum_{i'} B^{i'} f'_{i'} f'^2_{u'} - Q f'^3_{u'}, \end{aligned} \quad (4)$$

where $f'(t', x', u') = 0$ is the equation locally defining a submanifold L' of N' .

Recall that $F: (t, x, u) \mapsto (t', x', u')$ is a morphism in \mathcal{PE} if and only if for each point $\vartheta \in N$ and for each submanifold L' of N' , $F(\vartheta) \in L'$, the following two conditions are equivalent:

- the 2-jet of L' at the point $F(\vartheta)$ satisfies (4),
- the 2-jet of $F^{-1}(L')$ at the point ϑ satisfies (3).

In other words, the conditions “ f' is solution of (4)” and “ f is solution of (3)” should be equivalent when

$$f(t, x, u) = f'(t'(t, x, u), x'(t, x, u), u'(t, x, u)).$$

To find all such maps we use the following procedure:

1. Express derivatives of f in (3) through derivatives of f' :

$$\frac{\partial f}{\partial t} = \frac{\partial f'}{\partial t'} \frac{\partial t'}{\partial t} + \frac{\partial f'}{\partial x^{i'}} \frac{\partial x^{i'}}{\partial t} + \frac{\partial f'}{\partial u'} \frac{\partial u'}{\partial t}$$

and so on.

2. In the obtained identity substitute the combinations of the derivatives of f' for $\partial f'/\partial t'$ by formula (4). Then repeat this step for $\partial^2 f'/\partial t'^2$ in order to eliminate all derivatives with respect to t' . After reducing to common denominator, the transformed identity will have the form $\Phi = 0$, where Φ is a rational function of

partial derivatives of f' with respect to x' and u' . The coefficients ϕ_1, \dots, ϕ_s of Φ are functions of 4-jet of the map F .

3. Solve the system $\phi_1 = 0, \dots, \phi_s = 0$ of partial differential equations for a map F .

Let us realize this procedure. Note that we shall not write out function Φ completely. Instead we consider only some of its coefficients and then use the obtained information about F in order to simplify Φ step by step.

First note that the derivatives of the fourth order arise only in term $\partial^2 f' / \partial t'^2$ when we fulfill the step 2 of the above procedure. Write this term before the final realization of step 2 for the sake of simplicity:

$$\begin{aligned} \Phi &= \sum_{i,j} b^{ij} \left(t'_i t'_j f_u^2 - t'_i t'_u f_j f_u - t'_j t'_u f_i f_u + t_u^2 f_i f_j \right) \frac{\partial^2 f'}{\partial t'^2} + \dots \\ &= \sum_{i,j} b^{ij} (t'_i f_u - t'_u f_i) (t'_j f_u - t'_u f_j) \frac{\partial^2 f'}{\partial t'^2} + \dots \end{aligned}$$

The coefficient at $\partial^2 f' / \partial t'^2$ must be zero, and the quadratic form b^{ij} is positive definite. We get $t'_i f_u = t'_u f_i$, so

$$t'_i (f'_{t'} t'_u + f'_{x'} x'_u + f'_{u'} u'_u) = t'_u (f'_{t'} t'_i + f'_{x'} x'_i + f'_{u'} u'_i)$$

(here and below we use the notation

$$f'_{t'} = \frac{\partial f'}{\partial x'^{i'}}, \quad f'_{x'} x'_u = \sum_{j'} f'_{j'} x'^{j'}_u$$

and so on). Hence we obtain the following system of equations:

$$\begin{cases} t'_u u'_i = u'_u t'_i \\ t'_u x'_i = x'_u t'_i \end{cases} \quad (5)$$

One of the following three conditions holds:

1. $t'_u = 0, t'_x = 0$;
2. $t'_u = 0, t'_x \neq 0$;
3. $t'_u \neq 0$.

In the second case, $u'_u = x'_u = 0$. Taking into account the equality $t'_u = 0$, we obtain a desired contradiction to the assumption that F is a submersion.

In the third case, we get from (5) the identities $t'_x = \omega t'_u, u'_x = \omega u'_u, x'^{i'}_x = \omega x'^{i'}_u$, where $\omega = t'_x / t'_u$ is a section of $\pi^* T^* X$, $\pi: N = T \times X \times \Omega \rightarrow X$ is the natural projection, $\pi^* T^* X$ is the vector bundle over N induced by π from the cotangent bundle $T^* X$, and $\omega = \sum_i \omega_i(t, x, u) dx^i$ in local coordinates. This implies that $f_x = \omega f_u$. Substituting the last formula to (3), we get

$$f_t = f_u \left[\sum_{i,j} b^{ij} \left(\frac{\partial \omega_i}{\partial x^j} - \omega_j \frac{\partial \omega_i}{\partial u} \right) - \sum_{i,j} c^{ij} \omega_i \omega_j + \sum_i b^i \omega_i - q \right].$$

Denote the expression in square brackets by $\zeta(t, x, u)$. Then $f_t = \zeta f_u$. Expressing derivatives of f in terms of derivatives of f' , we obtain $t'_t = \zeta t'_u, x'_t = \zeta x'_u, u'_t = \zeta u'_u$. Consider the field of hyperplanes that kill the 1-form dt' in the tangent bundle TM (recall that $t'_u \neq 0$, so dt' is non-degenerated). The differential of F vanishes on these hyperplanes because $du' \wedge dt' = dx'^{i'} \wedge dt' = 0$. Therefore $\text{rank}(dF) \leq 1$. Since $\dim N' \geq 3$, F can not be submersive, which contradicts the definition of an admitted map.

Finally, we see that only the first case is possible. Hence t' is a function of t , and $f'_{t'}$ may appear only in the representation of f_t . Let us look at the terms of Φ containing $(f'_{u'})^{-2}$:

$$\Phi = \sum_{i',j',k',l'} t'_t x'^{i'}_u x'^{j'}_u B'^{k'l'} f'_{i'} f'_{j'} f'_{k'} f'_{l'} f'_{u'u'} (f'_{u'})^{-2} + \dots$$

Substitution of any covector $\omega = \sum_{i'} \omega_{i'} dx'^{i'}$ $\in \Gamma(T^* X')$ to the expression

$$\sum_{i',j',k',l'} t'_t x'^{i'}_u x'^{j'}_u B'^{k'l'} \omega_{i'} \omega_{j'} \omega_{k'} \omega_{l'} = t'_t \left(\sum_{i'} x'^{i'}_u \omega_{i'} \right)^2 \left(\sum_{k',l'} B'^{k'l'} \omega_{k'} \omega_{l'} \right)$$

should give zero. The quadratic form $B'^{k'l'}$ is positive definite, so $\sum_{k',l'} B'^{k'l'} \omega_{k'} \omega_{l'} > 0$ when $\omega \neq 0$. Taking into account that F is submersive, we obtain $t'_t \neq 0$. Therefore $\sum_{i'} x'^{i'} \omega_{i'} = 0$ for any ω , that is $x'^u \equiv 0$. This implies $x' = x'(t, x)$, $t' = t'(t)$, which completes the proof. \square

6 Comparison with the reduction by a symmetry group

As Remark 2 shows, our definition of morphism in \mathcal{PDE} is a generalization of the reduction by a symmetry group. So we can obtain sets of solutions more general than the sets of group-invariant solutions provided by the group analysis of PDE (though our approach is more laborious owing to the non-linearity of the system of PDE describing a morphisms). Let us illustrate this by an example of a primitive morphism.

Definition 10. A morphism $F: \mathbf{A} \rightarrow \mathbf{B}$ of a category \mathcal{C} is called a *reducible in \mathcal{C}* if there exist non-invertible morphisms $G: \mathbf{A} \rightarrow \mathbf{C}$, $H: \mathbf{C} \rightarrow \mathbf{B}$ in \mathcal{C} such that $F = H \circ G$. Otherwise, a morphism is called *primitive in \mathcal{C}* .

Note that the reduction of PDE by a symmetry group defines a primitive morphism if and only if this group has no proper subgroups. The reduction by any symmetry group that is not a discrete cyclic group of prime order may be always represented as a superposition of two nontrivial reductions, so the corresponding morphism is a superposition of two non-invertible morphisms and therefore is reducible. In particular, this situation takes place for any nontrivial connected Lie group.

However, the situation for morphisms is completely different. Even a morphism that decreases the number of independent variables by 2 or more may be primitive; below we present an example of such a morphism. In contrast, in the Lie group analysis we always have one-parameter subgroups of a symmetry group, so the morphism corresponding to a symmetry group is always reducible.

Example 3. Consider the following morphism $F: \mathbf{A} \rightarrow \mathbf{B}$ in \mathcal{PE} :

- \mathbf{A} is the heat equation $u_t = a(u)\Delta u$ posed on $X = \{(x, y, z, w) : z < w\} \subset \mathbb{R}^4$ equipped with the metric

$$g_{ij} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \gamma & \alpha & \beta \\ 0 & \alpha & 1 & 0 \\ 0 & \beta & 0 & 1 \end{pmatrix},$$

where $\alpha = xe^w$, $\beta = xe^z$, $\gamma = 1 + \alpha^2 + \beta^2$, $a \notin \mathcal{A}_{\text{exp}} \cup \mathcal{A}_{\text{deg}}$. In the coordinate form, \mathbf{A} looks as

$$\begin{aligned} a^{-1}(u)u_t &= u_{xx} + u_{yy} - 2\alpha u_{yz} - 2\beta u_{yw} + (1 + \alpha^2)u_{zz} \\ &+ 2\alpha\beta u_{zw} + (1 + \beta^2)u_{ww} + (\alpha\beta)_w u_z + (\alpha\beta)_z u_w. \end{aligned}$$

- \mathbf{B} is the heat equation $a^{-1}(u)u_t = u_{xx} + u_{yy}$ posed on $Y = \{(x, y)\} = \mathbb{R}^2$ equipped with Euclidean metric.
- The morphism F is defined by the map $(t, (x, y, z, w), u) \mapsto (t, (x, y), u)$.

This morphism decreases the number of independent variables by 2 and nevertheless is primitive in \mathcal{PE} .

Additional examples of morphisms that are not defined by any symmetry group of the given PDE, and also a detailed investigation of the case $\dim Y = \dim X - 1$, may be found in [6, sections 9, 10] and [7, sections 7–9].

Acknowledgements. This work was partially supported by the RFBR grants 15-01-02352 and 15-51-06001 (Russia). I am grateful to Vladimir Rubtsov and Yuri Zarhin for useful remarks.

References

- [1] V. I. Elkin. *Reduction of nonlinear control systems. A differential geometric approach*. Mathematics and its Applications, **472**. Kluwer Academic Publishers, Dordrecht, 1999.
- [2] J. Krasil'shchik, A. Vinogradov (eds.). *Symmetries and conservation laws for differential equations of mathematical physics*. Translations of Mathematical Monographs, **182**, AMS, Providence, RI, 1999.
- [3] S. Mac Lane. *Categories for the working mathematician*. Springer-Verlag, NY, 1998.
- [4] P. J. Olver. *Applications of Lie groups to differential equations*. Springer-Verlag, NY, 1993.

- [5] M. F. Prokhorova. *Modeling of solutions of heat equation and Stefan problem with dimension decrease.* Russian Acad. Sci. Docl. Math. **58** (1), 1998. 88–90.
- [6] M. F. Prokhorova. *Modeling of heat equation and Stefan problem.* Preprint VINITI N 347-V00, IMM UrBr Russian Acad. Sci., Ekaterinburg, 2000. 1–66 (in Russian).
- [7] M. F. Prokhorova. *Factorization of nonlinear heat equation posed on Riemann manifold.* arXiv:math/0108001 [math.AP], 2001. 1–9.
- [8] M. Prokhorova. *The structure of the category of parabolic equations.* arXiv:math/0512094v5 [math.AP], 2009, 1-33.
- [9] M. Prokhorova. *The structure of the category of parabolic equations. II.* Proceedings of the 47th International Youth School-conference “Modern Problems in Mathematics and its Applications”: 134-147, Yekaterinburg, Russia, 2016.