

Spectral Lattices of reducible matrices over completed idempotent semifields

Francisco J. Valverde-Albacete¹ and Carmen Peláez-Moreno^{2*}

¹ Departamento de Lenguajes y Sistemas Informáticos
Univ. Nacional de Educación a Distancia, c/ Juan del Rosal, 16. 28040 Madrid, Spain
`fva@lsi.uned.es`

² Departamento de Teoría de la Señal y de las Comunicaciones
Universidad Carlos III de Madrid, 28911 Leganés, Spain
`carmen@tsc.uc3m.es`

Abstract. Previous work has shown a relation between L-valued extensions of FCA and the spectra of some matrices related to L-valued contexts. We investigate the spectra of reducible matrices over completed idempotent semifields in the framework of dioids, naturally-ordered semirings, that encompass several of those extensions. Considering special sets of eigenvectors also brings out complete lattices in the picture and we argue that such structure may be more important than standard eigenspace structure for matrices over completed idempotent semifields.

1 Motivation

The eigenvectors and eigenspaces over certain naturally ordered semirings called *dioids* seem to be intimately related to multi-valued extensions of Formal Concept Analysis [1]. For instance [2, 3] prove that formal concepts are optimal factors for decomposing a matrix with entries in complete residuated semirings. Notice the strong analogy to the SVD, with formal concepts taking the role of pairs of left and right eigenvectors.

Indeed, [4] prove that, at least on a particular kind of dioids, the idempotent semifields, formal concepts are related to the eigenvectors of the unit in the semiring. This result, however, cannot be unified with the former both for theoretical reasons, since idempotent semifields are incomplete (see below), as well as for practical reasons, since the carrier set of idempotent semifields is almost never included in $[0, 1]$.

A possible way forward is to develop these theories under the common framework of the L -fuzzy sets, where L is any complete lattice [5]. Such an endeavour has already been called for [6], although it remains unfulfilled, hence this paper has a two-fold aim:

1. to clarify the spectral theory over *completed* idempotent semifields, and

* FJVA is supported by EU FP7 project LiMoSIne (contract 288024). CPM has been partially supported by the Spanish Government-Comisión Interministerial de Ciencia y Tecnología project TEC2011-26807 for this paper.

2. to take steps towards a common framework for the interpretation of *L-Formal Concept Analysis as a spectral construction*.

First steps have been taken in this direction with the development of a spectral theory of irreducible matrices [7] over complete idempotent semifields, whose summary we include below, but the general case, here presented, shows a more intimate relation to lattice theory.

1.1 Dioids and their spectral theory

A *semiring* is an algebra $\mathcal{S} = \langle S, \oplus, \otimes, \epsilon, e \rangle$ whose additive structure, $\langle S, \oplus, \epsilon \rangle$, is a commutative monoid and whose multiplicative structure, $\langle S \setminus \{\epsilon\}, \otimes, e \rangle$, is a monoid with multiplication distributing over addition from right and left and with additive neutral element absorbing for \otimes , i.e. $\forall a \in S, \epsilon \otimes a = \epsilon$.

Let $\mathcal{M}_n(\mathcal{S})$ be the semiring of square matrices over a semiring \mathcal{S} with the usual operations. Given $A \in \mathcal{M}_n(\mathcal{S})$ the *right (left) eigenproblem* is the task of finding the *right eigenvectors* $v \in S^{n \times 1}$ and *right eigenvalues* $\rho \in S$ (respectively *left eigenvectors* $u \in S^{1 \times n}$ and *left eigenvalues* $\lambda \in S$) satisfying:

$$u \otimes A = \lambda \otimes u \qquad A \otimes v = v \otimes \rho \qquad (1)$$

The left and right eigenspaces and spectra are the sets of these solutions:

$$\begin{aligned} \Lambda(A) &= \{\lambda \in S \mid \mathcal{U}_\lambda(A) \neq \{\epsilon^n\}\} & \text{P}(A) &= \{\rho \in S \mid \mathcal{V}_\rho(A) \neq \{\epsilon^n\}\} \\ \mathcal{U}_\lambda(A) &= \{u \in S^{1 \times n} \mid u \otimes A = \lambda \otimes u\} & \mathcal{V}_\rho(A) &= \{v \in S^{n \times 1} \mid A \otimes v = v \otimes \rho\} \\ \mathcal{U}(A) &= \bigcup_{\lambda \in \Lambda(A)} \mathcal{U}_\lambda(A) & \mathcal{V}(A) &= \bigcup_{\rho \in \text{P}(A)} \mathcal{V}_\rho(A) \end{aligned} \qquad (2)$$

Since $\Lambda(A) = \text{P}(A^T)$ and $\mathcal{U}_\lambda(A) = \mathcal{V}_\lambda(A^T)$, from now on we will omit references to left eigenvalues, eigenvectors and spectra, unless we want to emphasize differences.

With so little structure it might seem hard to solve (1), but a very generic solution based in the concept of transitive closure A^+ and transitive-reflexive closure A^* of a matrix is given by the following theorem:

Theorem 1 (Gondran and Minoux, [8, Theorem 1]). *Let $A \in S^{n \times n}$. If A^* exists, the following two conditions are equivalent:*

1. $A_{.i}^+ \otimes \mu = A_{.i}^* \otimes \mu$ for some $i \in \{1 \dots n\}$, and $\mu \in S$.
2. $A_{.i}^+ \otimes \mu$ (and $A_{.i}^* \otimes \mu$) is an eigenvector of A for e , $A_{.i}^+ \otimes \mu \in \mathcal{V}_e(A)$.

In [7] this result was made more specific in two directions: on the one hand, by focusing on particular types of completed idempotent semirings—semirings with a natural order where infinite additions of elements exist so transitive closures are guaranteed to exist and sets of generators can be found for the eigenspaces—and, on the other hand, by considering more easily visualizable subsemimodules than the whole eigenspace—a better choice for exploratory data analysis.

Specifically, every commutative semiring accepts a canonical preorder, $a \leq b$ if and only if there exists $c \in D$ with $a \oplus c = b$. A *dioid* is a semiring \mathcal{D}

where this relation is actually an order. Dioids are zerosumfree and entire, that is they have no non-null additive or multiplicative factors of zero. Commutative complete dioids are already complete residuated lattices. Similarly, semimodules over complete commutative dioids are also complete lattices.

An *idempotent semiring* is a dioid whose addition is idempotent, and a *selective semiring* one where the arguments attaining the value of the additive operation can be identified.

Example 1. Examples of idempotent dioids are

1. The *Boolean lattice* $\mathbb{B} = \langle \{0, 1\}, \vee, \wedge, 0, 1 \rangle$
2. All fuzzy semirings, e.g. $\langle [0, 1], \max, \min, 0, 1 \rangle$
3. The *min-plus algebra* $\mathbb{R}_{\min,+} = \langle \mathbb{R} \cup \{\infty\}, \min, +, \infty, 0 \rangle$
4. The *max-plus algebra* $\mathbb{R}_{\max,+} = \langle \mathbb{R} \cup \{-\infty\}, \max, +, -\infty, 0 \rangle$ □

Of the semirings above, only the boolean lattice and the fuzzy semirings are complete dioids, since the rest lack the *top* element \top as an adequate inverse for the bottom in the order.

The second important feature of spectra over complete dioids, as proven in [7], is that the set of eigenvalues on complete dioids is extended with respect to the incomplete case, and it makes sense to distinguish those which are associated to finite eigenvectors, the *proper eigenvalues* $P^P(A)$, and those associated with non-finite eigenvectors, the *improper eigenvalues* $P(A) \setminus P^P(A)$.

Moreover, as said above, the eigenspaces of matrices over complete dioids have the structure of a complete lattice. But since these lattices may be continuous and difficult to represent we introduce the more easily-represented *eigenlattices* $\mathcal{L}_\rho(A)$ and $\mathcal{L}_\lambda(A)$, complete *finite* sublattices of the eigenspaces to be used as scaffolding in visual representations.

1.2 Completed idempotent semifields and their spectral theory

A semiring is a *semifield* if there exists a multiplicative inverse for every element $a \in S$, notated as a^{-1} , and *radicable* if the equation $a^b = c$ can be solved for a . As exemplified above, idempotent semifields are incomplete in their natural order. Luckily, there are procedures for *completing* such structures [9] and we will not differentiate between *complete or completed* structures,

Example 2. The maxplus $\mathbb{R}_{\max,+}$ and minplus $\mathbb{R}_{\min,+}$ semifields can be completed as,

1. the *completed Minplus semifield*, $\overline{\mathbb{R}}_{\min,+} = \langle \mathbb{R} \cup \{-\infty, \infty\}, \min, \dot{+}, -\cdot, \infty, 0 \rangle$.
2. the *completed Maxplus semifield*, $\overline{\mathbb{R}}_{\max,+} = \langle \mathbb{R} \cup \{-\infty, \infty\}, \max, +, -\cdot, -\infty, 0 \rangle$,

In this notation we have $\forall c, -\infty + c = -\infty$ and $\infty \dot{+} c = \infty$, which solves several issues in dealing with the separately completed dioids. These two completions are inverses $\overline{\mathbb{R}}_{\min,+} = \overline{\mathbb{R}}_{\max,+}^{-1}$, hence order-dual lattices. Indeed they are better jointly called the *max-min-plus* semiring $\overline{\mathbb{R}}_{\max,+}^{\min,\dot{+}}$. □

In fact, idempotent semifields $\mathcal{K} = \langle K, \oplus, \dot{\oplus}, \otimes, \dot{\otimes}, \cdot^{-1}, \perp, e, \top \rangle$, appear as enriched structures, the advantage of working with them being that meets can be expressed by means of joins and inversion as $a \wedge b = (a^{-1} \oplus b^{-1})^{-1}$. On a practical note, residuation in complete commutative idempotent semifields can be expressed in terms of inverses, and this extends to eigenspaces.

Given $A \in \mathcal{M}_n(\mathcal{S})$, the *network (weighted digraph) induced by A*, $N_A = (V_A, E_A, w_A)$, consists of a set of vertices $V_A = \bar{n}$, a set of arcs, $E_A = \{(i, j) \mid A_{ij} \neq \epsilon_S\}$, and a weight $w_A : V_A \times V_A \rightarrow \mathcal{S}$, $(i, j) \mapsto w_A(i, j) = a_{ij}$. This allows us to apply intuitively all notions from networks to matrices and vice versa, like the underlying graph $G_A = (V_A, E_A)$, the set of paths $\Pi_A^+(i, j)$ between nodes i and j or the set of cycles C_A^+ . Matrix A is *irreducible* if every node of V_A is connected to every other node in V_A through a path, otherwise it is *reducible*.

We will use the spectra of irreducible matrices as a basic block for that of reducible matrices: if C_A^+ is the set of cycles of A then $\mu_{\oplus}(A) = \oplus\{\mu_{\oplus}(c) \mid c \in C_A^+\}$ is their *aggregate cycle mean*. For finite $\mu_{\oplus}(A)$, let $\tilde{A}^+ = (A \dot{\otimes} \mu_{\oplus}(A)^{-1})^+$ be the *normalized transitive closure of A*, and define the set of (right) *fundamental eigenvectors of irreducible A for ρ* as $\text{FEV}_{\rho}(A) = \{\tilde{A}_{\cdot i}^+ \mid \tilde{A}_{ii}^+ = e\}$, with left fundamental eigenvectors $\text{FEV}_{\rho}(A^T) = \text{FEV}_{\rho}(A)^T$. Then,

Theorem 2 ((Right) spectral theory for irreducible matrices, [7]). *Let $A \in \mathcal{M}_n(\overline{\mathcal{K}})$ be an irreducible matrix over a complete commutative selective radicable semifield. Then:*

1. $\Lambda(A) = \overline{\mathcal{K}} \setminus \{\perp\} = \text{P}(A)$.
2. $\Lambda^{\text{P}}(A) = \{\mu_{\oplus}(A)\} = \text{P}^{\text{P}}(A)$.
3. If $\rho \in \text{P}(A) \setminus \text{P}^{\text{P}}(A)$, then $\mathcal{V}_{\rho}(A) = \{\perp^n, \top^n\} = \mathcal{L}_{\rho}(A)$.
4. If $\rho = \mu_{\oplus}(A) < \top$, then $\mathcal{V}_{\rho}(A) = \langle \text{FEV}_{\rho}(A) \rangle_{\overline{\mathcal{K}}} \supset \mathcal{L}_{\rho}(A) = \langle \text{FEV}_{\rho}(A) \rangle_{\mathfrak{B}}$.

In this paper we try and find analogue results to Theorem 2 for the reducible case. First, we completely describe the spectra with Theorem 3 in Section 3.1. Then, we provide in Section 3.2 a bottom-up construction of the eigenspaces of certain reducible matrices from that of their irreducible blocks, using from Section 2.2 a recursive scheme to render matrices over idempotent semifields into specialised Upper Frobenius Normal Forms (UFNF). Finally, we discuss our findings in Section 4 and relate them to previous approaches.

2 Preliminaries

2.1 Some partial orders and lattices

In this paper we assume familiarity with basic order notions as described in [1, 10]. We only introduce notation when departing from there.

Recall that every set V with $|V| = n$ elements and a total order $\leq \subseteq V \times V$ is isomorphic to a lattice called the *chain of n elements*, $\langle V, \leq \rangle \cong \mathbf{n}$. When the relation is the empty order relation $\emptyset \in V \times V$, it is called an *anti-chain of n elements*, $\langle V, \emptyset \rangle \cong \bar{\mathbf{n}}$. Lattice $\mathbb{1} \cong \mathbf{1}$ is the vacuously-ordered singleton. Lattice

2 is the boolean lattice isomorphic to chain **2**. Lattice **3** is a lattice lying at the heart of completed semifields, the **3**-bounded lattice-ordered group $\perp < e < \top$, isomorphic to chain **3**.

If set of order ideals of a poset P is $\mathcal{O}(P)$, then

Proposition 1 ([10, Chap. 1]). *Let $\langle P, \leq \rangle$ be a finite poset. Then $\langle \mathcal{O}(P), \subseteq \rangle$ is a lattice obtained by the embedding $\varphi : P \rightarrow \mathcal{O}(P), \varphi(x) = \downarrow x$, with $\forall A_1, A_2 \in \mathcal{O}(P), A_1 \vee A_2 = A_1 \cup A_2$ and $A_1 \wedge A_2 = A_1 \cap A_2$.*

Note that $x \leq y$ in \mathcal{P} if and only if $\downarrow x \subseteq \downarrow y$ in $\mathcal{O}(P)$. Furthermore, $\mathcal{O}(P)$ can be obtained systematically from the ordered set in a number of cases:

Proposition 2 ([10, Chap. 1]). *Let $\langle P, \leq \rangle$ be a finite poset. Then*

1. $\mathcal{O}(P \oplus \mathbb{1}) \cong \mathcal{O}(P) \oplus \mathbb{1}$ and $\mathcal{O}(\mathbb{1} \oplus P) \cong \mathbb{1} \oplus \mathcal{O}(P)$.
2. $\mathcal{O}(P_1 \uplus P_2) \cong \mathcal{O}(P_1) \times \mathcal{O}(P_2)$.
3. $\mathcal{O}(P^d) \cong \mathcal{F}(P) \cong \mathcal{O}(P)^d$.
4. $\mathcal{O}(n) \cong n \oplus \mathbb{1} \cong \mathbb{1} \oplus n$.
5. $\mathcal{O}(\bar{n}) \cong 2^n$.

2.2 An inductive structure for reducible matrices

Recall that a *digraph* (or *directed graph*), is a pair $G = (V, E)$ with V a set of *vertices* and $E \subseteq V \times V$ a set of *arcs* (*directed edges*), ordered pairs of vertices, such that for every $i, j \in V$ there is at most one arc $(i, j) \in E$. If $(i, j) \in E$ then we say that “ i is a predecessor of j ” or “ j is a successor of i ”, and $E \in \mathcal{M}_n(\mathbb{B})$ is the *associated relation* of G . If there is a walk from a vertex i to a vertex j in G we say that “ i has access to j ” or j is *reachable from i* , $i \rightsquigarrow j$. Hence, *reachability* is the transitive closure of the associated relation, $\rightsquigarrow = E^+$ [11]. However, vertices $j \in V$ with no incoming or outgoing arcs cannot be part of any cycle, hence $j \not\rightsquigarrow j$ for such nodes, so it is not reflexive, in general. $(\rightsquigarrow \cap I_V)$ is the *reflexive restriction* of \rightsquigarrow , that is, the biggest reflexive relation included in it.

If there is a walk from a vertex i to vertex j in G or viceversa we say that i and j are *connected*, $i \rightsquigarrow j \vee j \rightsquigarrow i$. *Connectivity* is the symmetric closure of the reachability relation: its transitive, reflexive restriction is an equivalence relation on V_G whose classes are called the *(dis)connected components* of G . Note that each of the (outwards) disconnected components is actually (inwards) connected. Let $K \geq 1$ be the number of disconnected components of G . Then V and E are partitioned into the subsets of vertices $\{V_k\}_{k=1}^K$ and arcs $\{E_k\}_{k=1}^K$ of each disconnected component $\bigcup_k V_k = V$, $V_k \cap V_l = \emptyset, k \neq l$, $\bigcup_k E_k = E$, $E_k \cap E_l = \emptyset, k \neq l$ and we may write $G = \uplus_{k=1}^K G_k$ is a disjoint union of graphs. G is called *connected* itself if $K = 1$.

On the other hand, since reachability is transitive by construction, its symmetric, reflexive restriction $i \rightsquigarrow j \Leftrightarrow i \rightsquigarrow j \wedge j \rightsquigarrow i$ is a refinement of connectivity called *strong connectivity*. Its equivalence classes are the *strongly connected components* of G . For each disconnected component G_k , let R_k be the number of its strongly connected components. If $R_k = 1$ then the k -th component is *strongly*

connected, otherwise just *connected* and composed of a number of strongly connected components itself. G is *strongly connected* itself if $K = R = 1$.

Given a digraph $G = (V, E)$, the *reduced or condensation digraph*, $\overline{G} = (\overline{V}, \overline{E})$ is induced by the set $\overline{V} = V / \rightsquigarrow$ of strongly connected components, and $C, C' \in \overline{V}$, $(C, C') \in \overline{E}$ iff there exists one arc $(i, j) \in E$ for some $i \in C, j \in C'$ and we say that component C has access to C' , and write $C \preccurlyeq C'$. It is well known that $\overline{G} = (\overline{V}, \overline{E})$ is a partially ordered set called a *directed acyclic graph (dag)*.

Given a matrix $A \in \mathcal{M}_n(S)$, its *condensation digraph* is the partial order of strong connectivity classes $\overline{G}_A = (\overline{V}_A, \overline{E}_A)$, as in Figure 2b in Example 4, of its associated digraph $G_A = (V_A, E_A)$. This can be proven by means of an Upper Frobenius Normal Form (UFNF) [12], a structure for matrices that we intend to specialise and use as a scheme for structural induction over reducible matrices.

In the following, for a set of indices $V_x \subseteq \overline{\mathbf{n}}$ we write $P(V_x)$ for a permutation of it, and \mathcal{E}_{xy} is an empty matrix of conformal dimension most of the times represented on matrix patterns as elliptical dots.

Lemma 1 (Recursive Upper Frobenius Normal Form, UFNF). *Let $A \in \mathcal{M}_n(S)$ be a matrix over a semiring and \overline{G}_A its condensation digraph. Then,*

1. (UFNF₃) *If A has zero lines it can be transformed by a simultaneous row and column permutation of V_A into the following form:*

$$P_3^T \otimes A \otimes P_3 = \begin{bmatrix} \mathcal{E}_{\iota\iota} & \cdot & \cdot & \cdot \\ \cdot & \mathcal{E}_{\alpha\alpha} & A_{\alpha\beta} & A_{\alpha\omega} \\ \cdot & \cdot & A_{\beta\beta} & A_{\beta\omega} \\ \cdot & \cdot & \cdot & \mathcal{E}_{\omega\omega} \end{bmatrix} \quad (3)$$

where: either $A_{\alpha\beta}$ or $A_{\alpha\omega}$ or both are non-zero, and either $A_{\alpha\omega}$ or $A_{\beta\omega}$ or both are non-zero. Furthermore, P_3 is obtained concatenating permutations for the indices of simultaneously zero columns and rows V_ι , the indices of zero columns but non-zero rows V_α , the indices of zero rows but non-zero columns V_ω and the rest V_β as $P_3 = P(V_\iota)P(V_\alpha)P(V_\beta)P(V_\omega)$.

2. (UFNF₂) *If A has no zero lines it can be transformed by a simultaneous row and column permutation $P_2 = P(A_1) \dots P(A_k)$ into block diagonal UFNF*

$$P_2^T \otimes A \otimes P_2 = \begin{bmatrix} A_1 & \cdot & \dots & \cdot \\ \cdot & A_2 & \dots & \cdot \\ \vdots & \vdots & \ddots & \vdots \\ \cdot & \cdot & \dots & A_K \end{bmatrix} \quad (4)$$

where $\{A_k\}_{k=1}^K, K \geq 1$ are the matrices of connected components of \overline{G}_A .

3. (UFNF₁) *If A is reducible with no zero lines and a single connected component it can be simultaneously row- and column-permuted by P_1 to*

$$P_1^T \otimes A \otimes P_1 = \begin{bmatrix} A_{11} & A_{12} & \dots & A_{1R} \\ \cdot & A_{22} & \dots & A_{2R} \\ \vdots & \vdots & \ddots & \vdots \\ \cdot & \cdot & \dots & A_{RR} \end{bmatrix} \quad (5)$$

where A_{rr} are the matrices associated to each of its R strongly connected components (sorted in a topological ordering), and $P_1 = P(A_{11}) \dots P(A_{RR})$.

The particular choice of UFNF is clarified by the following Lemma, since the condensation digraph will prove important later on:

Lemma 2 (Scheme for structural induction over reducible matrices). Let $A \in \mathcal{M}_n(S)$ be a matrix over an entire zerosumfree semiring and \overline{G}_A its condensation digraph. Then:

1. If A is irreducible then $\overline{G}_A \cong \mathbb{1}$.
2. If A is in UFNF₂ then $\overline{G}_A = \bigsqcup \overline{G}_{A_k}$.
3. If A is in UFNF₃ then $\overline{G}_A = \overline{G}_{A_{\beta\beta}}$.
4. $\overline{G}_{A^r} = (\overline{G}_A)^d$.

Note that for A in UFNF₁, \overline{G}_A may be any connected ordered set.

3 Results

3.1 Generic results for reducible matrices

The following lemma clarifies the order relation between eigenvectors in ordered semimodules,

Lemma 3. Let \mathcal{X} be a naturally-ordered semimodule.

1. Vectors with incomparable supports are incomparable.
2. If \mathcal{X} is further complete, vectors with incomparable saturated supports are incomparable.

Proof. Let v and w be two vectors in \mathcal{X} . Comparability of supports lies in the \subseteq relation: if $\text{supp}(v) \parallel \text{supp}(w)$ then $\text{supp}(v) \not\subseteq \text{supp}(w)$ and $\text{supp}(w) \not\subseteq \text{supp}(v)$. Therefore from $\text{supp}(v) \cap \text{supp}(w)^C \neq \emptyset$ we have $v(\text{supp}(v) \cap \text{supp}(w)^C) \neq \perp$ and $w(\text{supp}(v) \cap \text{supp}(w)^C) = \perp$, hence $v \not\leq w$. Similarly, from $\text{supp}(w) \cap \text{supp}(v)^C \neq \emptyset$ we have $w \not\leq v$, therefore $v \parallel w$. Claim 2 is likewise argued on the support taking the role of $\overline{\mathbf{n}}$, and the saturated support taking the role of the original support. \square

Let $A \in \mathcal{M}_n(\mathcal{S})$ be a matrix and \overline{G}_A be its condensation digraph. Consider a class $C_r \in \overline{V}_A$ and call $V_u = (\bigcup_{C' \in \downarrow C_r} C') \setminus C_r$, $V_d = (\bigcup_{C' \in \uparrow C_r} C') \setminus C_r$ and $V_p = V_A \setminus (V_u \cup C_r \cup V_d)$ the sets of upstream, downstream and parallel vertices for C_r , respectively. Using permutation $P_r = P(V_u)P(C_r)P(V_p)P(V_d)$ we may suppose a blocked form of $A(C_r)$ like the one in Fig. 1 without loss of generality. Notice that any of V_u, V_d or V_p may be empty. However, if V_u (resp. V_d) is not of null dimension, then A_{ur} (resp. A_{rd}) cannot be empty.

Proposition 3. Let $A \in \mathcal{M}_n(\overline{\mathcal{K}})$ be a matrix over a complete commutative selective radicable semifield with $C_A^+ \neq \emptyset$. Then a scalar $\rho > \perp$ is a proper eigenvalue of A if and only if there is at least one class in its condensation digraph $C_r \in \overline{G}_A$ such that $\rho = \mu_{\oplus}(A_{rr})$.




Fig. 1: Matrix A focused on C_r , $A(C_r) = P_r^T \otimes A \otimes P_r$ and associated digraph. The loops at each node, consisting of (possibly empty) A_{uu} , A_{rr} , A_{pp} , A_{dd} are not shown.

Proof. The proof, for instance, in [8] extends even to $\rho = \top$,. □

The proper spectrum is clarified by:

Lemma 4. *Let $A \in \mathcal{M}_n(\mathcal{S})$ be a reducible matrix over a complete radicable selective semifield. Then, there are no other finite eigenvectors in $\mathcal{V}_\rho(A)$ contributed by \tilde{A}^ρ than those selected by the critical circuits in $C_r \in \bar{V}_A$ such that $\mu_\oplus(A_{rr}) = \rho$,*

$$\text{FEV}^F(A) = \cup_{C_r \in \bar{V}_A} \{(\tilde{A}^\rho)_{\cdot i}^+ \mid i \in V_r^c, \mu_\oplus(A_{rr}) = \rho\} .$$

Proof. If $\rho = \mu_\oplus(A_{rr})$, from Proposition 3 we see that the finite eigenvectors mentioned really belong in $\mathcal{V}_\rho(A)$. If $\rho > \mu_\oplus(A_{rr})$ then $(\tilde{A}^\rho)_{ii}^+ < e = (\tilde{A}^\rho)_{ii}^*$ hence the columns selected by C_r do not generate eigenvectors. If $\rho < \mu_\oplus(A_{rr})$ then $(\tilde{A}^\rho)_{ij}^+ = \top$ and whether those classes with cycle mean ρ are upstream or downstream of C_r the only value that is propagated is \top , hence the eigenvectors are all saturated. □

Theorem 3 (Spectra of generic matrices). *Let $A \in \mathcal{M}_n(\bar{\mathcal{D}})$ be a reducible matrix over an entire zerosumfree semiring. Then,*

1. *If $C_A^+ = \emptyset$ then $P(A) = P^P(A) = \{\epsilon\}$.*
2. *If $C_A^+ \neq \emptyset$ and further $\bar{\mathcal{D}}$ is a complete selective radicable semifield,*
 - (a) *If $\bar{zc}(A) \neq \emptyset$ then $P(A) = \bar{\mathcal{K}}$ and $P^P(A) = \{\perp\} \cup \{\mu_\oplus(A_{rr}) \mid C_r \in \bar{V}_A\}$.*
 - (b) *If $\bar{zc}(A) = \emptyset$ then $P(A) = \bar{\mathcal{K}} \setminus \{\perp\}$ and $P^P(A) = \{\mu_\oplus(A_{rr}) \mid C_r \in \bar{V}_A\}$.*

Proof. First, call $\bar{zc}(A)$ the set of empty columns of A . If G_A has no cycles $C_A^+ = \emptyset$, claim 1 follows from a result in [7]. But if $C_A^+ \neq \emptyset$ then by Proposition 3, $P^P(A) \supseteq \{\mu_\oplus(A_{rr}) \mid C_r \in \bar{V}_A\}$ and no other non-null proper eigenvalues may exist by Lemma 4. \perp is only proper when $\bar{zc}(A) \neq \emptyset$ hence claims 2a and 2b follow. □

Translating to UFNF terms:

Corollary 1. *Let $A \in \mathcal{M}_n(\overline{\mathcal{K}})$ be a matrix over a complete selective radicable semifield with $C_A^+ \neq \emptyset$. Then $P(A) = \overline{\mathcal{K}} \setminus \{\perp\}$ and $P^p(A) = \{\mu_{\oplus}(A_{rr}) \mid C_r \in \overline{V}_A\}$, unless A is in $UFNF_3$ and $\overline{zc}(A) \neq \emptyset$ whence $\perp \in P^p(A) \subseteq P(A)$ too.*

Proof. If A is irreducible or in $UFNF_1$ or $UFNF_2$ it has no empty columns or rows. This can only happen in $UFNF_3$ in which case the result follows from Theorem 3. \square

Since this solves entirely the description of the spectrum, only the description of the eigenspaces is left pending.

3.2 Eigenspaces of matrices in $UFNF_1$

In this section we offer an instance of how the UFNF can be used to obtain, inductively, the spectrum of reducible matrices.

If for every parallel condensation class $C_p \subseteq V_A$ in $A(C_r)$ illustrated in C_r of Fig. 1 $A_{up} \neq \mathcal{E}_{up}$ or $A_{pd} \neq \mathcal{E}_{pd}$ or both, then A is in $UFNF_1$ with a single connected block. In this case, we can relate the order of the eigenvectors to the condensation digraph: define the *support of a class* $\text{supp}(C)$ as the support of any of the non-null eigenvectors it induces in A .

Lemma 5. *Let $A \in \mathcal{M}_n(\mathcal{S})$ be a matrix in $UFNF_1$ over a complete zerosumfree semiring. Then, for any class $C_r \in \overline{V}_A$, $\text{supp}(C_r) = \bigcup \{C_{l_r} \mid C_{l_r} \in \downarrow C_r\}$.*

Proof. Since A_{rr} is irreducible, if $\rho = \mu_{\oplus}(A_{rr})$ then for any $v_r \in \mathcal{V}_{\rho}(A_{rr})$ we have that $\text{supp}(v_r) = V_r$, hence $V_r \subseteq \text{supp}(C_r)$. Also, since \mathcal{S} is complete and zerosumfree $(\tilde{A}^{\rho})_{rr}^+$ exists and is full [7]. Since $(\tilde{A}^{\rho})_{uu}^+ \tilde{A}_{ur}^{\rho}$ must have a full column for any $C_{l_r} \in \downarrow C_r$ meaning precisely that C_r is downstream from C_{l_r} , the product $(\tilde{A}^{\rho})_{uu}^+ \tilde{A}_{ur}^{\rho} (\tilde{A}^{\rho})_{rr}^+$ for the nodes in C_{l_r} is non-null and $V_{l_r} \subseteq \text{supp}(C_r)$. \square

Lemma 5 establishes a bijection between the supports of condensation classes and downsets in \overline{G}_A which is actually an isomorphism of orders,

Proposition 4. *Let $A \in \mathcal{M}_n(\overline{\mathcal{K}})$ be a matrix over a commutative complete selective radicable semifield admitting an $UFNF_1$. Then*

1. *Each class $C_r \in \overline{V}_A$ generates a distinct saturated eigenvector, v_r^{\top} .*
2. *$\{v_r^{\top} \mid C_r \in \overline{V}_A\} \cong \overline{G}_A$.*

Proof. Let $v \in \mathcal{V}_{\rho}(A)$ where $\rho = \mu_{\oplus}(A_{rr})$ then by Lemma 5 $\text{supp}(v) = \downarrow C_r$, hence $v_r^{\top} = \top v \in \mathcal{V}_{\rho}(A)$ is the unique saturated eigenvector, since $\text{sat-supp}(\top v) = \text{supp}(\top v) = \text{supp}(C)$, and the bijection follows. By Lemma 3, claim 2 the order relation between classes is maintained between eigenvectors, whence the order isomorphism in claim 2. \square

We call $\text{FEV}^{1,\top}(A) = \{v_r^\top \mid C_r \in \overline{V}_A\}$ the *set of saturated fundamental eigenvectors of A*. Notice that $\overline{V}_{A^\top} = \overline{V}_A$ but $\overline{E}_{A^\top} = \overline{E}_A^d$, so $\text{FEV}^{1,\top}(A^\top) \cong \overline{G}_A^d$.

For every finite $\rho \in \text{P}^{\text{P}}(A)$ we define the *critical nodes* $V_\rho^c = \{i \in \overline{\mathbf{n}} \mid (\tilde{A}^\rho)_{ii}^+ = e\}$ and $\text{FEV}_\rho^{1,\text{F}}(A) = \{(\tilde{A}^\rho)_{\cdot i}^+ \mid i \in V_\rho^c\}$ the *(maybe partially) finite fundamental eigenvectors of ρ* . Next, let $\delta_\rho^{-1}(\rho') = e$ if $\rho' = \rho$ and $\delta_\rho^{-1}(\rho') = \top$ otherwise. for $\rho \in \text{P}(A)$ the set of *(right) fundamental eigenvectors of A in UFNF₁ for ρ* as

$$\text{FEV}_\rho^1(A) = \cup_{\rho' \in \text{P}(A)} \{\delta_\rho^{-1}(\rho') \otimes \text{FEV}_{\rho'}^{1,\text{F}}(A)\} . \tag{6}$$

This definition absorbs two cases, actually,

Lemma 6. *Let $A \in \mathcal{M}_n(\overline{\mathcal{K}})$ be a matrix over a commutative complete selective radicable semifield admitting an UFNF₁. Then,*

1. *for $\rho \in \text{P}(A) \setminus \text{P}^{\text{P}}(A)$, $\text{FEV}_\rho^1(A) = \text{FEV}^{1,\top}(A)$.*
2. *for $\rho \in \text{P}^{\text{P}}(A)$, $\rho \neq \top$, $\text{FEV}_\rho^1(A) = \text{FEV}_\rho^{1,\text{F}}(A) \cup \text{FEV}^{1,\top}(A) \setminus (\top \otimes \text{FEV}_\rho^{1,\text{F}}(A))$.*
3. *for $\rho \in \text{P}(A)$, $\rho \neq \top$, $\text{FEV}^{1,\top}(A) = \top \otimes \text{FEV}_\rho^1(A)$.*

Proof. If $\rho \in \text{P}(A) \setminus \text{P}^{\text{P}}(A)$, then for all $\rho' \in \overline{\mathcal{K}}$, $\delta_\rho^{-1}(\rho') = \top$. By Proposition 4 claim 1 follows as we range $\rho' \in \text{P}^{\text{P}}(A)$. Similarly, when $\rho \in \text{P}^{\text{P}}(A)$, those classes whose $\rho' \neq \rho$ supply a single saturated eigenvector. However, if $\rho' = \rho$, then $\delta_\rho^{-1}(\rho') = e$ obtains the (partially) finite fundamental eigenvectors $\text{FEV}_\rho^{1,\text{F}}(A)$, the saturated eigenvectors of which cannot be considered fundamental, since they can be obtained from $\text{FEV}_{\rho'}^{1,\text{F}}(A)$ linearly, and will not appear in $\text{FEV}_\rho^1(A)$. Claim 3 is a corollary of the other two. \square

Call $\mathcal{V}^\top(A) = \langle \text{FEV}^{1,\top}(A) \rangle_{\overline{\mathcal{K}}}$ the *saturated eigenspace of A*, then

Corollary 2. *Let $A \in \mathcal{M}_n(\overline{\mathcal{K}})$ be a matrix over a commutative complete selective radicable semifield admitting an UFNF₁. Then,*

1. *For $\rho \in \text{P}(A)$, $\mathcal{V}^\top(A) \subseteq \mathcal{V}_\rho(A)$.*
2. *For $\rho \in \text{P}(A) \setminus \text{P}^{\text{P}}(A)$, furthermore, $\mathcal{V}^\top(A) = \mathcal{V}_\rho(A)$.*

Proof. Since we have $\text{FEV}^{1,\top}(A) \subseteq \mathcal{V}_\rho(A)$, then $\mathcal{V}^\top(A) \subseteq \mathcal{V}_\rho(A)$. For $\rho \in \text{P}(A) \setminus \text{P}^{\text{P}}(A)$, $\text{FEV}_\rho^1(A) = \text{FEV}^{1,\top}(A)$ by Lemma 6, so claim 2 follows. \square

Hence, $\mathcal{V}^\top(A)$ provides a common “scaffolding” for every eigenspace, while the peculiarities for proper eigenvalues are due to the finite eigenvectors. Also, since $\mathcal{V}^\top(A)$ is a complete lattice, $\text{FEV}^{1,\top}(A) \subseteq \mathcal{V}^\top(A)$ is actually a lattice embedding,

Proposition 5. *Let $A \in \mathcal{M}_n(\overline{\mathcal{K}})$ be a matrix over a commutative complete selective radicable semifield admitting an UFNF₁. Then*

1. *For $\rho \in \text{P}(A) \setminus \text{P}^{\text{P}}(A)$,*

$$\mathcal{U}^\top(A) = \langle \text{FEV}^{1,\top}(A^\top) \rangle_{\mathfrak{F}} \cong \mathcal{F}(\overline{G}_A) \quad \mathcal{V}^\top(A) = \langle \text{FEV}^{1,\top}(A) \rangle_{\mathfrak{F}} \cong \mathcal{O}(\overline{G}_A) . \tag{7}$$

2. for all $\rho \in P^P(A)$, $\rho < \top$

$$\mathcal{U}_\lambda(A) = \langle \text{FEV}_\lambda^1(A^\top)^\top \rangle_{\overline{\mathcal{K}}} \quad \mathcal{V}_\rho(A) = \langle \text{FEV}_\rho^1(A) \rangle_{\overline{\mathcal{K}}} .$$

Proof. If $v_r^\top \in \text{FEV}^{1,\top}(A)$ then $\lambda v_r^\top = \lambda(\top v_r^\top) = v_r^\top$, whence $\mathcal{V}^\top(A) = \langle \text{FEV}^{1,\top}(A) \rangle_{\mathfrak{B}}$. In fact, the generation process may proceed on only a subsemiring of $\overline{\mathcal{K}}$ which need not even be complete. For instance, we may use any of the isomorphic copies of 2 embedded in $\overline{\mathcal{K}}$, for instance $\{\perp, k\} \cong 2$, with $k \neq \perp$.

Since the number of saturated eigenvectors is finite, being identical to the number of condensation classes, we only have to worry about binary joins and meets. Recall that $v_r^\top \vee v_k^\top = v_r^\top \oplus v_k^\top$ and $v_r^\top \wedge v_k^\top = v_r^\top \dot{\oplus} v_k^\top = \left((v_r^\top)^{-1} \dot{\oplus} (v_k^\top)^{-1} \right)^{-1}$. Then $\text{supp}(v_r^\top \oplus v_k^\top) = \text{supp}(v_r^\top) \cup \text{supp}(v_k^\top)$ and also

$$\text{supp}(v_r^\top \dot{\oplus} v_k^\top) = (\text{supp}^c(v_r^\top) \cup \text{supp}^c(v_k^\top))^c = \text{supp}(v_r^\top) \cap \text{supp}(v_k^\top)$$

for $C_r, C_k \in \overline{V}_A$ and Proposition 1 gives the result. For $\rho \in P^P(A)$, $\text{FEV}_\rho^1(A) \subseteq \mathcal{V}_\rho(A)$ implies that $\langle \text{FEV}_\rho^1(A) \rangle_{\overline{\mathcal{K}}} \subseteq \mathcal{V}_\rho(A)$, and Corollary 4 ensures that no finite vectors are missing. And dually for left eigenspaces. \square

This actually proves that $\text{FEV}_\rho^1(A)$ is join-dense in $\mathcal{V}_\rho(A)$.

As already mentioned, $\mathcal{V}_\rho(A)$ is a hard-to-visualize semimodule. An *eigen-space schematics* is a modified order diagram where the saturated eigenspace is represented in full but the rays generated by finite eigenvalues $\{\kappa \otimes (\tilde{A}^\rho)_{\cdot i}^\top \mid i \in V_r^c, \rho = \mu_{\oplus}(A_{rr})\}$ are drawn with discontinuous lines, as in the examples below.

We are now introducing another representation inspired by (7): we define the (left) right eigenlattices of A for $(\lambda \in \Lambda(A)) \rho \in P(A)$ as


$$\mathcal{L}_\lambda(A) = \langle \text{FEV}_\rho^1(A^\top)^\top \rangle_{\mathfrak{B}} \quad \mathcal{L}_\rho(A) = \langle \text{FEV}_\rho^1(A) \rangle_{\mathfrak{B}} .$$

Example 3 (Spectral lattices of irreducible matrices). Since irreducible matrices are in UFNF_1 with a single class, $\text{FEV}_{\mu_{\oplus}(A)}^0(A) = \text{FEV}_{\mu_{\oplus}(A)}^1(A)$. For $\rho \in P(A) \setminus P^P(A)$ we have $\text{FEV}^{0,\top}(A) = \{\top^n\}$, whence $\overline{G}_A \cong \mathbb{1}$ and $\mathcal{V}^\top(A) = \{\perp^n, \top^n\} \cong 2$. For $\rho \in P^P(A)$, $\rho < \top$, as proven in [7], $\mathcal{V}_\rho(A)$ is finitely generable from $\text{FEV}_\rho^0(A)$, but the form of the eigenspace and eigenlattice for $A^P(A) = \{\mu_{\oplus}(A)\} = P^P(A)$ depends on the critical cycles and the eigenvectors they induce. \square

Example 4. Consider the matrix $A \in \mathcal{M}_n(\overline{\mathbb{R}}_{\max,+})$ from [13, p. 25.7, example 2] in UFNF_1 depicted in Fig. 2.(a). The condensed graph \overline{G}_A in Fig. 2.(b) has for vertex set $\overline{V}_A = \{C_1 = \{1\}, C_2 = \{2, 3, 4\}, C_3 = \{5, 6, 7\}, C_4 = \{8\}\}$, so consider the strongly connected components $G_{A_{kk}} = (C_k, E \cap C_k \times C_k)$, $1 \leq k \leq 4$. Their maximal cycle means are $\mu_k = \mu_{\oplus}(A_{kk})$: $\mu_1 = 0$, $\mu_2 = 2$, $\mu_3 = 1$ and $\mu_4 = -3$, respectively, corresponding to critical circuits: $C^c(G_{A_{11}}) = \{1 \circlearrowleft\}$, $C^c(G_{A_{22}}) = \{2 \rightarrow 3 \rightarrow 2\}$, $C^c(G_{A_{33}}) = \{5 \circlearrowleft, 6 \rightarrow 7 \rightarrow 6\}$, $C^c(G_{A_{44}}) = \{8 \circlearrowleft\}$.

$$A_3 = \begin{bmatrix} 0 & 0 & 7 & \dots & \dots & \dots & \dots & \dots \\ \dots & 3 & 0 & \dots & \dots & \dots & \dots & \dots \\ \dots & 1 & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & 2 & \dots & \dots & \dots & \dots & 10 & \dots \\ \dots & \dots & 1 & 0 & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & 0 & \dots & \dots & \dots \\ \dots & \dots & \dots & -1 & 2 & \dots & 23 & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & -3 \end{bmatrix}$$

(a) A reducible matrix in $UFNF_1$




(b) Class diagram (rectangles) overlaid on G_{A_3}

$$\begin{matrix} 1 \\ 2 \\ 3 \\ 5 \\ 6 \\ 7 \\ 8 \end{matrix} \begin{bmatrix} 0 & \top & \top & \top & \top & \top & \top & \top \\ \cdot & 0 & 1 & -2 & \cdot & \cdot & \cdot & 6 \\ \cdot & -1 & 0 & -3 & \cdot & \cdot & \cdot & 5 \\ \cdot & \cdot & \cdot & \cdot & 0 & -1 & -2 & 20 \\ \cdot & \cdot & \cdot & \cdot & -3 & 0 & -1 & 21 \\ \cdot & \cdot & \cdot & \cdot & -2 & 1 & 0 & 22 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 0 \end{bmatrix}$$


(c) Left fundamental eigenvectors

$$\begin{matrix} 1 & 2 & 3 & 5 & 6 & 7 & 8 \end{matrix} \begin{bmatrix} 0 & -3 & -2 & 6 & 5 & 4 & \top \\ \cdot & 0 & 1 & \cdot & \cdot & \cdot & \top \\ \cdot & -1 & 0 & \cdot & \cdot & \cdot & \top \\ \cdot & 0 & 1 & \cdot & \cdot & \cdot & \top \\ \cdot & \cdot & \cdot & 0 & -1 & -2 & \top \\ \cdot & \cdot & \cdot & -3 & 0 & -1 & \top \\ \cdot & \cdot & \cdot & -2 & 1 & 0 & \top \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 0 \end{bmatrix}$$


(d) Right fundamental eigenvectors



(e) $\mathcal{V}^\top(A_3)$



(f) Schematics of $\mathcal{V}_2(A_3)$



(g) Schematics of $\mathcal{V}(A_3)$

Fig. 2: Matrix A_3 (a), its associated digraph and class diagram (b), its left (c) and right (d) fundamental eigenvectors annotated with their eigennodes to the left and above, respectively; the eigenspace of improper eigenvectors $\mathcal{V}^\top(A_3)$ in (e), a schematic of the right eigenspace of proper eigenvalue $\rho = 2$, $\mathcal{V}_2(A_3)$ in (f) and the schematics of the whole right eigenspace $\mathcal{V}(A_3)$ in (g).

Note that node 4 does not generate an eigenvector in either spectrum, since it does not belong to a critical cycle.

Therefore $P(A_3) = P^P(A_3) = \{2, 1, 0, -3\}$ each left eigenspace is the span of the set of eigenvectors chosen from distinct critical cycles for each class of A : $\mathcal{U}_{\mu_1}(A) = \langle (\tilde{A}_3^{\mu_1})_{1.}^+ \rangle$, $\mathcal{U}_{\mu_2}(A) = \langle (\tilde{A}_3^{\mu_2})_{2.}^+ \rangle$, $\mathcal{U}_{\mu_3}(A) = \langle (\tilde{A}_3^{\mu_3})_{\{5,6\}.}^+ \rangle$, and $\mathcal{U}_{\mu_4}(A) = \langle (\tilde{A}_3^{\mu_4})_{8.}^+ \rangle$ —as described by the row vectors of Fig. 2.(c)—and the right eigenspaces are $\mathcal{V}_{\mu_1}(A) = \langle (\tilde{A}_3^{\mu_1})_{.1}^+ \rangle$, $\mathcal{V}_{\mu_2}(A) = \langle (\tilde{A}_3^{\mu_2})_{.2}^+ \rangle$, $\mathcal{V}_{\mu_3}(A) = \langle (\tilde{A}_3^{\mu_3})_{. \{5,6\}}^+ \rangle$, and $\mathcal{V}_{\mu_4}(A) = \langle (\tilde{A}_3^{\mu_4})_{.8}^+ \rangle$ —as described by the column vectors of Fig. 2.(d).

The saturated eigenspace is easily represented by means of an order diagram—Hasse diagram—as that of Fig. 2.(e). Note how it is embedded in that of any proper eigenvalue like $\rho = 2$ in Fig. 2.(f). Since the representation of continuous eigenspaces is problematic, we draw *schematics* of them, as in Fig. 2.(f). Fig. 2.(g) shows a schematic view of the union of the eigenspaces for proper eigenvalues $\mathcal{V}(A_3) = \cup_{\rho \in P^P(A)} \mathcal{V}_{\rho}(A_3)$. \square

4 Discussion

In this paper, we have discussed the spectrum of reducible matrices with entries in completed idempotent semifields. To the extent of our knowledge, this was pioneered in [14] and both [7] and this paper can be understood as systematic explorations to try and understand what was stated in there. For this purpose, the consideration of particular UFNF forms for the matrices has been crucial: while the description in [14] is combinatorial ours is constructive.

The usual notion of spectrum as the set of eigenvectors with more than one (null) eigenvector appears in this context as too weak: when a matrix has at least one cycle then all the values in the semifield (except the bottom \perp) belong to the spectrum. If the matrix has at least one empty column (resp. empty row) and a cycle then *all* of the semifield is the spectrum. Rather than redefine the notion of spectrum we have decided to introduce the *proper spectrum* as the set of eigenvalues with at least one vector with finite support.

Regarding the eigenspaces, we found not only that they are complete continuous lattices for proper eigenvalues, but also that they are *finite (complete) lattices* for *improper* eigenvalues. Looking for a device to represent the information within each proper eigenspace we focus on the fundamental eigenvectors of an irreducible matrix for each eigenvalue: those with unit values in certain of their coordinates. The span of those eigenvectors by the action of the \mathfrak{B} -bounded lattice-ordered group generates the finite eigenlattices. Interestingly, since improper eigenvectors only have non-finite coordinates, their span by the \mathfrak{B} -blog is exactly the same finite lattice as their span by the whole semifield itself.

With these building blocks it is easy to build finite lattices for reducible matrices of any UFNF description, as exemplified above. We believe this will

be a useful technique to understand and visualize the concept lattices of formal contexts with entries in an idempotent semifield and other dioids.

Bibliography

- [1] Ganter, B., Wille, R.: Formal Concept Analysis: Mathematical Foundations. Springer, Berlin, Heidelberg (1999)
- [2] Belohlavek, R., Vychodil, V.: Discovery of optimal factors in binary data via a novel method of matrix decomposition. *Journal Of Computer And System Sciences* **76**(1) (2010) 3–20
- [3] Belohlavek, R.: Optimal decompositions of matrices with entries from residuated lattices. *Journal Of Logic And Computation* **22**(6) (November 2012) 1405–1425
- [4] Valverde-Albacete, F.J., Pelaez-Moreno, C.: Spectral lattices of $(R_{max,+})$ -formal contexts. In: Formal Concept Analysis. Volume LCNS 4933. (2008) 124–139
- [5] Goguen, J.A.: L-fuzzy sets. *J. Math. Anal. Appl* **18**(1) (1967) 145–174
- [6] Gondran, M., Minoux, M.: Dioids and semirings: links to fuzzy set and other applications. *Fuzzy Sets And Systems* **158**(1273–1294) (April 2007)
- [7] Valverde-Albacete, F.J., Peláez-Moreno, C.: The spectra of irreducible matrices over completed commutative idempotent semifields. In preparation (2013)
- [8] Gondran, M., Minoux, M.: Valeurs propres et vecteurs propres dans les dioïdes et leur interprétation en théorie des graphes. EDF, Bulletin de la Direction des Etudes et Recherches, Serie C, Mathématiques Informatique **2** (1977) 25–41
- [9] Valverde-Albacete, F.J., Pelaez-Moreno, C.: Extending conceptualisation modes for generalised Formal Concept Analysis. *Information Sciences* **181** (May 2011) 1888–1909
- [10] Davey, B., Priestley, H.: Introduction to lattices and order. 2nd edn. Cambridge University Press, Cambridge, UK (2002)
- [11] Schmidt, G., Ströhlein, T.: Relations and Graphs. Discrete Mathematics for Computer Scientists. Springer (1993)
- [12] Brualdi, R.A., Ryser, H.J.: Combinatorial Matrix Theory. Volume 39 of Encyclopedia of Mathematics and its Applications. Cambridge University Press (1991)
- [13] Akian, M., Bapat, R., Gaubert, S.: 25. [15] 25–1–25–17
- [14] Jun, M., Yan, G., zhi Yong, D.: The eigen-problem in the completely max-algebra. In: Proceedings of the Sixth International Conference on Parallel and Distributed Computing. (2005)
- [15] Hogben, L., ed.: Handbook of Linear Algebra. Discrete Mathematics and Its Applications. Chapman & Hall/CRC (2007)