

Interpolant Existence is Undecidable for Two-Variable First-Order Logic with Two Equivalence Relations

Frank Wolter¹, Michael Zakharyashev²

¹University of Liverpool, Ashton Street, Liverpool L69 3BX, UK

³Birkbeck, University of London, Malet Street, London WC1E 7HX, UK

Abstract

The interpolant existence problem (IEP) for a logic L is to decide, given formulas φ and ψ , whether there exists a formula ι , built from the shared symbols of φ and ψ , such that φ entails ι and ι entails ψ in L . If L enjoys the Craig interpolation property (CIP), then the IEP reduces to validity in L . Recently, the IEP has been studied for logics without the CIP. The results obtained so far indicate that even though the IEP can be computationally harder than validity, it is decidable when L is decidable. Here, we give the first examples of decidable fragments of first-order logic for which the IEP is undecidable. Namely, we show that the IEP is undecidable for the two-variable fragment with two equivalence relations and for the two-variable guarded fragment with individual constants and two equivalence relations. We also determine the corresponding decidable Boolean description logics for which the IEP is undecidable.

Keywords

Craig interpolant, interpolant existence problem, two-variable fragment of first-order logic.


1. Introduction

A first-order (FO) formula ι is called a Craig interpolant for FO-formulas φ and ψ if $\varphi \models \iota \models \psi$ and ι is built from the shared non-logical symbols of φ and ψ . Interpolants have been applied in many areas ranging from formal verification [1] and software specification [2] to theory combinations [3, 4, 5] and query reformulation and rewriting in databases [6, 7].


For many fragments L of FO, including many description logics (DLs), the existence of a Craig interpolant for φ and ψ is guaranteed if φ entails ψ . This phenomenon is known as the Craig interpolation property (CIP) of L . The CIP holds, for instance, for the DLs \mathcal{ALC} , \mathcal{ALCT} , and \mathcal{ALCQI} [8], and also for the guarded negation fragment of FO [9]. In this case, the problem to decide whether an interpolant for φ and ψ exists reduces to showing that φ entails ψ in L , and so is not harder than entailment. Moreover, interpolants can often be extracted from a proof of the entailment.

An important consequence of the CIP of a logic L is the projective Beth definability property (BDP) of L : if a relation is implicitly definable over a signature ρ in L , then it is explicitly definable over ρ in L . The BDP is used in ontology engineering for extracting explicit definitions of concepts or nominals from ontologies [10, 8, 11], developing and maintaining ontology alignments [12], and for robust modularisations and decompositions of ontologies [13, 14]. Explicit definitions are used in ontology-based data management to equivalently rewrite ontology-mediated queries [15, 16, 17, 18, 19].

Unfortunately, there are many prominent fragments of FO that do not enjoy the CIP and BDP: for example, DLs with nominals such as \mathcal{ALCO} and/or role inclusions such as \mathcal{ALCH} [8], the guarded and the two-variable fragment of FO [20, 21, 22]. To extend interpolation-based techniques to such formalisms, it has recently been suggested to investigate the interpolant existence problem (IEP) for logics L without the CIP: given formulas φ and ψ as an input, decide whether they have a Craig interpolant in L . It has been shown that the IEP is decidable for many DLs with nominals and role inclusions [23], Horn-DLs [24], the two-variable and guarded fragments of FO [25], and some decidable

 DL 2024: 37th International Workshop on Description Logics, June 18–21, 2024, Bergen, Norway

 wolter@liverpool.ac.uk (F. Wolter); m.zakharyashev@bbk.ac.uk (M. Zakharyashev)

 0000-0002-4470-606X (F. Wolter); 0000-0002-2210-5183 (M. Zakharyashev)



© 2024 Copyright for this paper by its authors. Use permitted under Creative Commons License Attribution 4.0 International (CC BY 4.0).

fragments of first-order modal logics [26]. These positive results naturally lead to a conjecture that the IEP for L is decidable whenever the entailment in L is decidable.

In this paper, we disprove this conjecture by showing that the IEP is undecidable for the two-variable FO with two equivalence relations and also for the two-variable guarded fragment of FO with constants and two equivalence relations. In the realm of Description Logic, the former result means that interpolant existence is undecidable for concept inclusions in \mathcal{ALC} enriched with Boolean operators on roles and the identity role. As a consequence, we also obtain the undecidability of the explicit definition existence problem (EDEP) for each of these logics L , which is the problem of deciding, given formulas φ, ψ and a signature ϱ , whether there exists an explicit ϱ -definition of ψ modulo φ in L . (It is known [26] that the IEP and EDEP are polynomially reducible to each other.)

The paper is structured as follows. Section 2 introduces the fragments of FO we are interested in and summarises the necessary technical tools and results. Section 3 proves the undecidability of the IEP for the two-variable FO with two equivalence relations by reduction of the infinite Post Correspondence Problem. Section 4 shows how this result can be adapted to the two-variable guarded fragment with constants and two equivalence relations and to suitable description logics. Finally, Section 5 discusses a few challenging open problems.

2. Preliminaries

Let σ be a signature of individual constants and *unary and binary* predicate symbols, including two distinguished binary *equivalence predicates* E_1, E_2 . Let var be a set comprising two individual variables. We consider the following fragments of first-order logic $\text{FO}(\sigma)$ over σ :

$\text{FO}^2\text{2E}(\sigma)$ is the set of *constant-free* FO-formulas constructed in the usual way from atoms $x = y$ with $x, y \in var$ and $R(\mathbf{x})$, for any n -ary predicate symbol $R \in \sigma$ and n -tuple \mathbf{x} of variables from var ;

$\text{GF}^2\text{2E}_c(\sigma)$ admits variables in var and individual constants in atoms but restricts quantification to the patterns

$$\forall \mathbf{y} (\alpha(\mathbf{x}, \mathbf{y}) \rightarrow \varphi(\mathbf{x}, \mathbf{y})), \quad \exists \mathbf{y} (\alpha(\mathbf{x}, \mathbf{y}) \wedge \varphi(\mathbf{x}, \mathbf{y})),$$

where $\varphi(\mathbf{x}, \mathbf{y})$ is a $\text{GF}^2\text{2E}_c$ -formula and $\alpha(\mathbf{x}, \mathbf{y})$ is an atom with occurrences of \mathbf{x}, \mathbf{y} .

The *signature* $\text{sig}(\psi)$ of a formula ψ in any of these logics is the set of predicate and constant symbols occurring in ψ . Formulas are interpreted in σ -structures $\mathfrak{A} = (\text{dom}(\mathfrak{A}), (R^{\mathfrak{A}})_{R \in \sigma}, (c^{\mathfrak{A}})_{c \in \sigma})$ with a domain $\text{dom}(\mathfrak{A}) \neq \emptyset$, relations $R^{\mathfrak{A}}$ on $\text{dom}(\mathfrak{A})$ of the same arity as predicates $R \in \sigma$, and $c^{\mathfrak{A}} \in \text{dom}(\mathfrak{A})$. It is always assumed that $E_1^{\mathfrak{A}}$ and $E_2^{\mathfrak{A}}$ are *equivalence relations* on $\text{dom}(\mathfrak{A})$. A *pointed structure* is a pair \mathfrak{A}, \mathbf{a} with a tuple $\mathbf{a} = (a_1, \dots, a_n)$, $n \leq 2$, of elements from $\text{dom}(\mathfrak{A})$.

The decision problems for $\text{FO}^2\text{2E}$ and $\text{GF}^2\text{2E}_c$ are known to be co2NEXP TIME - and 2EXP TIME -complete, respectively; see [27, 28] and references therein.

Definition 1. Let $L \in \{\text{FO}^2\text{2E}, \text{GF}^2\text{2E}_c\}$, let $\varphi(\mathbf{x})$ and $\psi(\mathbf{x})$ be $L(\sigma)$ -formulas with the same free variables \mathbf{x} , and let $\varrho = \text{sig}(\varphi) \cap \text{sig}(\psi)$. An $L(\varrho)$ -formula $\iota(\mathbf{x})$ is called an *L -interpolant* for φ and ψ if $\varphi(\mathbf{x}) \models \iota(\mathbf{x})$ and $\iota(\mathbf{x}) \models \psi(\mathbf{x})$.

Our concern here is the following *L -interpolant existence problem* (L -IEP, for short):

(L -IEP) given L -formulas $\varphi(\mathbf{x})$ and $\psi(\mathbf{x})$, decide whether they have an L -interpolant $\iota(\mathbf{x})$.

We remind the reader of a well-known criterion of interpolant existence in terms of appropriate bisimulations. Let $\varrho \subseteq \sigma$. Given pointed σ -structures \mathfrak{A}, \mathbf{a} and \mathfrak{B}, \mathbf{b} , we call \mathfrak{A}, \mathbf{a} and \mathfrak{B}, \mathbf{b} *$L(\varrho)$ -equivalent* and write $\mathfrak{A}, \mathbf{a} \equiv_{L, \varrho} \mathfrak{B}, \mathbf{b}$ if $\mathfrak{A} \models \varphi(\mathbf{a})$ iff $\mathfrak{B} \models \varphi(\mathbf{b})$, for all $L(\varrho)$ -formulas φ .

Definition 2. A nonempty relation $\beta \subseteq \text{dom}(\mathfrak{A}) \times \text{dom}(\mathfrak{B})$ is called an *$\text{FO}^2\text{2E}(\varrho)$ -bisimulation between \mathfrak{A} and \mathfrak{B}* if the following conditions are satisfied for all $(a, b) \in \beta$:

1. for every $a' \in \text{dom}(\mathfrak{A})$, there is a $b' \in \text{dom}(\mathfrak{B})$ such that $(a', b') \in \beta$ and $(a, a') \mapsto (b, b')$ is a partial ϱ -isomorphism between \mathfrak{A} and \mathfrak{B} ;
2. for every $b' \in \text{dom}(\mathfrak{B})$, there is a $a' \in \text{dom}(\mathfrak{A})$ such that $(a', b') \in \beta$ and $(a, a') \mapsto (b, b')$ is a partial ϱ -isomorphism between \mathfrak{A} and \mathfrak{B} .

Observe that $\text{FO}^2\text{2E}(\varrho)$ -bisimulations are *global* in the sense that $\text{dom}(\mathfrak{A}) \subseteq \{a \mid (a, b) \in \beta\}$ and $\text{dom}(\mathfrak{B}) \subseteq \{b \mid (a, b) \in \beta\}$.

Definition 3. A global relation $\beta \subseteq \text{dom}(\mathfrak{A}) \times \text{dom}(\mathfrak{B})$ is an $\text{GF}^2\text{2E}_c(\varrho)$ -bisimulation between \mathfrak{A} and \mathfrak{B} if $(c^{\mathfrak{A}}, c^{\mathfrak{B}}) \in \beta$ for all $c \in \varrho$ and the following conditions are satisfied for all $(a, b) \in \beta$:

1. for every $a' \in \text{dom}(\mathfrak{A})$ such that either $R^{\mathfrak{A}}(a, a')$, for some binary $R \in \varrho$, or $a' = a$, condition 1 of Definition 2 holds;
2. for every $b' \in \text{dom}(\mathfrak{B})$ such that either $R^{\mathfrak{B}}(b, b')$, for some binary $R \in \varrho$, or $b' = b$, condition 2 of Definition 2 holds.

For tuples $\mathbf{a} = (a_1, \dots, a_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$ with $n \leq 2$, we write $\mathfrak{A}, \mathbf{a} \sim_{L, \varrho} \mathfrak{B}, \mathbf{b}$ if $\mathbf{a} \mapsto \mathbf{b}$ is a partial ϱ -isomorphism between \mathfrak{A} and \mathfrak{B} and there is an $L(\varrho)$ -bisimulation β between \mathfrak{A} and \mathfrak{B} such that $(a_i, b_i) \in \beta$, for all $i \leq n$. The following characterisation is well known; see, e.g., [29, 30, 28]:

Lemma 1. Let $L \in \{\text{FO}^2\text{2E}, \text{GF}^2\text{2E}_c\}$ and $\varrho \subseteq \sigma$. For any pointed σ -structures \mathfrak{A}, \mathbf{a} and \mathfrak{B}, \mathbf{b} with $|\mathbf{a}| = |\mathbf{b}|$,

$$\mathfrak{A}, \mathbf{a} \sim_{L, \varrho} \mathfrak{B}, \mathbf{b} \quad \text{implies} \quad \mathfrak{A}, \mathbf{a} \equiv_{L, \varrho} \mathfrak{B}, \mathbf{b}$$

and, conversely, if structures \mathfrak{A} and \mathfrak{B} are ω -saturated [31], then

$$\mathfrak{A}, \mathbf{a} \equiv_{L, \varrho} \mathfrak{B}, \mathbf{b} \quad \text{implies} \quad \mathfrak{A}, \mathbf{a} \sim_{L, \varrho} \mathfrak{B}, \mathbf{b}.$$

Using this lemma, one can obtain the following variant of Robinson's joint consistency criterion [32, 25]:

Lemma 2. Let $L \in \{\text{FO}^2\text{2E}, \text{GF}^2\text{2E}_c\}$. Then $L(\sigma)$ -formulas $\varphi(\mathbf{x})$ and $\psi(\mathbf{x})$ do not have an L -interpolant iff there exist pointed $L(\sigma)$ -structures \mathfrak{A}, \mathbf{a} and \mathfrak{B}, \mathbf{b} such that

- $\mathfrak{A} \models \varphi(\mathbf{a})$,
- $\mathfrak{B} \models \neg\psi(\mathbf{b})$,
- $\mathfrak{A}, \mathbf{a} \sim_{L, \varrho} \mathfrak{B}, \mathbf{b}$, where $\varrho = \text{sig}(\varphi) \cap \text{sig}(\psi)$.

Definition 4. Given formulas $\varphi(\mathbf{x})$, $\psi(\mathbf{x})$ and a signature ϱ , an *explicit ϱ -definition of $\psi(\mathbf{x})$ modulo $\varphi(\mathbf{x})$ in a logic L* is an $L(\varrho)$ -formula $\chi(\mathbf{x})$ such that $\models \varphi(\mathbf{x}) \rightarrow (\psi(\mathbf{x}) \leftrightarrow \chi(\mathbf{x}))$.

The *explicit ϱ -definition existence problem (EDEP)* for L is formulated as follows:

(L-EDEP) given L -formulas $\varphi(\mathbf{x})$, $\psi(\mathbf{x})$ and a signature ϱ , decide whether there exists an explicit ϱ -definition of $\psi(\mathbf{x})$ modulo $\varphi(\mathbf{x})$ in L .

The IEP and EDEP turn out to be closely related [33]. Here, we only need the following lemma whose proof can be found in [26]:

Lemma 3. For any $L \in \{\text{FO}^2\text{2E}, \text{GF}^2\text{2E}_c\}$, the L -IEP is polynomially reducible to the L -EDEP.

3. Undecidability of the FO²2E-IEP

In this section, we prove the undecidability of the FO²2E-IEP by a reduction of the undecidable *infinite Post Correspondence Problem* (ω PCP) [34, 35], which is formulated as follows: given an alphabet $\Gamma = \{A_1, \dots, A_n\}$, $n \geq 2$, and a finite set \mathcal{P} of pairs $(v_1, w_1), \dots, (v_k, w_k)$ of non-empty words over Γ , decide whether there exists an infinite sequence of indices i_1, i_2, \dots , for $1 \leq i_j \leq k$ and $j < \omega$, such that the ω -words $v_{i_1}v_{i_2}\dots$ and $w_{i_1}w_{i_2}\dots$ coincide. If this is the case, the ω PCP instance \mathcal{P} is said to *have a solution*.

Suppose that an ω PCP instance \mathcal{P} is given. Our aim is to construct FO²2E-formulas $\varphi(x)$ and $\psi(x)$ such that \mathcal{P} has a solution iff $\varphi(x)$ and $\neg\psi(x)$ are satisfied in FO²2E(ρ)-bisimilar pointed structures, where ρ is the shared signature of φ and ψ , and then apply Lemma 2. In our construction $\rho = \{R, S\} \cup \Gamma$ with two binary predicates R, S and the $A_i \in \Gamma$ treated as unary predicates. The formula $\varphi(x)$ is defined by taking:

$$\begin{aligned} \varphi(x) = & X_0(x) \wedge \exists y (R(x, y) \wedge X_1(y)) \wedge \forall x [X_1(x) \rightarrow \exists y (R(x, y) \wedge X_2(y))] \wedge \\ & \forall x [X_2(x) \rightarrow \exists y (R(x, y) \wedge X_0(y))] \wedge \forall x, y ((X_0(x) \wedge X_0(y)) \rightarrow x = y) \wedge \\ & \exists y [S(x, y) \wedge \bigwedge_{A_i \in \Gamma} (A_i(y) \rightarrow A_i(x))]. \end{aligned}$$

The only purpose of $\varphi(x)$ is to generate an R -cycle with three points: for any structure \mathfrak{A} , a_0 , if $\mathfrak{A} \models \varphi(a_0)$, then \mathfrak{A} contains a cycle $R(a_0, a_1), R(a_1, a_2), R(a_2, a_0)$. The last conjunct of φ is needed to ensure that $\text{sig}(\varphi) \cap \text{sig}(\psi) = \rho$.

Before defining $\psi(x)$ formally, we explain the intuition behind $\neg\psi(x)$. Suppose \mathfrak{A}, a_0 and \mathfrak{B}, b_0 are such that $\mathfrak{A} \models \varphi(a_0)$, $\mathfrak{B} \models \neg\psi(b_0)$ and $\mathfrak{A}, a_0 \sim_{L, \rho} \mathfrak{B}, b_0$, for $L = \text{FO}^2\text{2E}$. Then \mathfrak{B} contains an R -chain $\dots b_{-3}, b_{-2}, b_{-1}, b_0, b_1, b_2, b_3, \dots$ such that $\mathfrak{A}, a_0 \sim_{L, \rho} \mathfrak{B}, b_{3m}$, for $m \in \mathbb{Z}$. The purpose of the formula $\neg\psi(x)$ is to generate an infinite S -chain \bar{v} starting from b_0 , along which an infinite sequence of words $v_i, i < \omega$, is written, and also an infinite S -chain \bar{w} starting from b_3 , along which a sequence of words $w_i, i < \omega$, is written. Using the equivalence relations E_1 and E_2 , the formula $\neg\psi(x)$ ensures that the respective pairs $(v_i, w_i), i < \omega$, are all in \mathcal{P} . That the resulting ω -words coincide (and give a solution to \mathcal{P}) is ensured by $\mathfrak{A}, a_0 \sim_{L, \rho} \mathfrak{B}, b_0$. The intended models \mathfrak{A}, a_0 and \mathfrak{B}, b_0 are illustrated in Fig. 1.

To define $\neg\psi$ formally, suppose $v_j = A_1^j \dots A_{n_j}^j$ and $w_j = B_1^j \dots B_{m_j}^j$, for $1 \leq j \leq k$, $n_j \geq 1$, and $m_j \geq 1$. Along with the $A_i \in \Gamma$, to generate unique S -chains \bar{v} and \bar{w} , we require auxiliary unary predicates $P_{v_j}^l, l = 1, \dots, n_j$, and $P_{w_j}^l, l = 1, \dots, m_j$, as well as Z_i^v and Z_i^w , for $i = 1, 2$, and Z^v, Z^w . Let $\bar{1} = 2, \bar{2} = 1, \bar{x} = y$, and $\bar{y} = x$.

We define $\neg\psi(x)$ to be a conjunction of the following four groups of formulas:

$$\begin{aligned} \text{(generation)} \quad & Z_1^v(x) \wedge Z^v(x), \quad \forall x, y (Z^u(x) \wedge S(x, y) \rightarrow Z^u(y)), \quad \text{for } u \in \{v, w\}, \\ & \forall x (Z_i^v(x) \rightarrow \bigvee_{j \leq k} \nu_i^j(x)), \quad \forall x (Z_i^w(x) \rightarrow \bigvee_{j \leq k} \omega_i^j(x)), \quad \text{for } i = 1, 2, \\ & \forall y (R(x, y) \rightarrow Y_1(y)), \quad \forall x, y (Y_1(x) \wedge R(x, y) \rightarrow Y_2(y)), \\ & \quad \quad \quad \forall x, y (Y_2(x) \wedge R(x, y) \rightarrow Z_1^w(y) \wedge Z^w(y)), \end{aligned}$$

where, the formula ν_i^j generating the word v_j , for $i = 1, 2$, is defined recursively as

$$\begin{aligned} \nu_i^j(x) &= \alpha_{i,1}^j(x) \wedge \beta_{i,1}^j(x), \\ \alpha_{i,l}^j(x) &= \exists \bar{x} (S(x, \bar{x}) \wedge E_i(x, \bar{x}) \wedge A_l^j(\bar{x}) \wedge P_{v_j}^l(\bar{x}) \wedge \alpha_{i,l+1}^j(\bar{x})), \quad \text{for } l = 1, \dots, n_j - 1, \\ \alpha_{i,n_j}^j(x) &= \exists \bar{x} (S(x, \bar{x}) \wedge E_i(x, \bar{x}) \wedge A_{n_j}^j(\bar{x}) \wedge P_{v_j}^{n_j}(\bar{x}) \wedge Z_i^v(\bar{x})), \\ \beta_{i,l}^j(x) &= \forall \bar{x} (S(x, \bar{x}) \rightarrow E_i(x, \bar{x}) \wedge A_l^j(\bar{x}) \wedge P_{v_j}^l(\bar{x}) \wedge \beta_{i,l+1}^j(\bar{x})), \quad \text{for } l = 1, \dots, n_j - 1, \\ \beta_{i,n_j}^j(x) &= \forall \bar{x} (S(x, \bar{x}) \rightarrow E_i(x, \bar{x}) \wedge A_{n_j}^j(\bar{x}) \wedge P_{v_j}^{n_j}(\bar{x}) \wedge Z_i^v(\bar{x})), \end{aligned}$$




Figure 1: Intended models \mathfrak{A}, a_0 and \mathfrak{B}, b_0 with $\mathfrak{A} \models \varphi(a_0)$, $\mathfrak{B} \models \neg\psi(b_0)$, and $\mathfrak{A}, a_0 \sim_{L, \varrho} \mathfrak{B}, b_0$, where ϱ -bisimilar points in the disjoint union of \mathfrak{A} and \mathfrak{B} are connected by dotted lines and E_i -equivalence classes in \mathfrak{B} , $i = 1, 2$, are encircled by dashed lines. Model \mathfrak{A} contains two disjoint copies of a 3-point R -cycle followed by an infinite S -chain. Model \mathfrak{B} has an infinite R -chain of points b_i , for $i \in \mathbb{Z}$; each b_{6i} starts an infinite S -chain for the word $v_{i_1} v_{i_2} \dots$, and each b_{6i+3} starts an infinite S -chain for the word $w_{i_1} w_{i_2} \dots$. These ω -words coincide with the ω -word written on the S -chains in \mathfrak{A} .

and the formulas ω_i^j , generating the respective words w_j , are defined analogously but with $B_l^j, P_{w_j}^l$ and m_j in place of $A_l^j, P_{v_j}^l$ and n_j , respectively.

(disjointness) $\forall x (P(x) \rightarrow \neg P'(x))$, for any distinct $P, P' \in \Gamma$, distinct P, P' of the form $P_{v_j}^l, P_{w_j}^l$, as well as for $\{P, P'\} = \{Z_1^u, Z_2^u\}$ with $u \in \{v, w\}$, and $\{P, P'\} = \{Z^v, Z^w\}$.

(coordination) $\forall x, y (R(x, y) \rightarrow E_1(x, y))$,
 $\forall x [Z_i^v(x) \wedge \nu_i^j(x) \rightarrow \forall y (E_i(x, y) \wedge Z_i^w(y) \rightarrow \omega_i^j(y))]$, for $i = 1, 2, j \leq k$,
 $\forall x, y (E_i(x, y) \wedge Z_i^v(x) \wedge Z_i^w(y) \rightarrow E_i(x, y))$, for $i = 1, 2$.

(uniqueness) $\forall x, y (P(x) \wedge P(y) \wedge E_i(x, y) \rightarrow x = y)$, for P of the form $P_{v_j}^l, P_{w_j}^l$, $i = 1, 2$.


Clearly $\varrho = \text{sig}(\varphi) \cap \text{sig}(\psi)$ consists of R, S and the predicates in Γ .

We now show that the constructed FO²2E-formulas φ and ψ are as required:

Theorem 4. Let φ and ψ be the FO^2E -formulas defined above for a given ωPCP instance \mathcal{P} . Then \mathcal{P} has a solution iff there exist structures \mathfrak{A}, \mathbf{a} and \mathfrak{B}, \mathbf{b} such that $\mathfrak{A} \models \varphi(\mathbf{a})$, $\mathfrak{B} \models \neg\psi(\mathbf{b})$, and $\mathfrak{A}, \mathbf{a} \sim_{\text{FO}^2\text{E}, \varrho} \mathfrak{B}, \mathbf{b}$.

Proof. (\Rightarrow) Suppose that $v_{i_1}v_{i_2}\dots = w_{i_1}w_{i_2}\dots$ is a solution to the given \mathcal{P} . It is not hard to check that, for the structures \mathfrak{A} and \mathfrak{B} shown in Fig. 1, we have $\mathfrak{A} \models \varphi(a_0)$, $\mathfrak{B} \models \neg\psi(b_0)$, and $\mathfrak{A}, a_0 \sim_{\text{FO}^2\text{E}(\varrho)} \mathfrak{B}, b_0$. It is to be noted that a formula enforcing a 3-point cycle is needed as a 1- or 2-point cycle is not $\text{FO}^2\text{E}(\varrho)$ -bisimilar to a chain. Also note that two disjoint copies of a 3-point R -cycle followed by an S -chain in \mathfrak{A} are needed to ensure $\text{FO}^2\text{E}(\varrho)$ -bisimilarity between \mathfrak{A}, a_0 and \mathfrak{B}, b_0 . Indeed, suppose, for example, that a and b are the i th points in the S -chains starting from a_0 and b_0 , respectively. Now, if we consider the $(i+1)$ th point b' in the S -chain starting from b_3 , then, to obtain a partial ϱ -isomorphism between (b, b') and some (a, a') , we may need to take the $(i+1)$ th point a' from the second S -chain in \mathfrak{A} .

(\Leftarrow) Suppose $\mathfrak{A} \models \varphi(a_0)$, $\mathfrak{B} \models \neg\psi(b_0)$, and $\mathfrak{A}, a_0 \sim_{\text{FO}^2\text{E}, \varrho} \mathfrak{B}, b_0$. Consider first the pointed structure \mathfrak{B}, b_0 . As $\mathfrak{B} \models \neg\psi(b_0)$, the axioms in the first two lines of **(generation)** give an infinite sequence $\bar{v} = \bar{v}_{i_1}\bar{v}_{i_2}\dots$ depicted below.



We show that the ω -word $v_{i_1}v_{i_2}\dots$ over Γ , i.e.,

$$\underbrace{A_1^{i_1} A_2^{i_1} \dots A_{n_{i_1}}^{i_1}}_{v_{i_1}} \underbrace{A_1^{i_2} A_2^{i_2} \dots A_{n_{i_2}}^{i_2}}_{v_{i_2}} \dots$$

written on \bar{v} is unique. Indeed, by **(disjointness)** and **(generation)**, all S -successors of b_0 carry the same $P_{v_{i_1}}^1$, all of them are in the same E_1 -equivalence class, and so must coincide by **(uniqueness)**. Next we see that all of the S -successors of the $P_{v_{i_1}}^1$ -point carry $P_{v_{i_1}}^2$, and so must coincide, etc. This gives us the word v_{i_1} carried by the segment \bar{v}_{i_1} sitting in the same E_1 -equivalence class. The last point in this segment carries Z_2^v , which triggers the next segment \bar{v}_{i_2} sitting in the same E_2 -equivalence class, and so on. All of the points in \bar{v} carry Z^v by the second formula in **(generation)**. The points on the boundaries between \bar{v}_{i_j} and $\bar{v}_{i_{j+1}}$ belong to both E_1 - and E_2 -equivalence classes.

As $\mathfrak{A}, a_0 \sim_{L, \varrho} \mathfrak{B}, b_0$, $R \in \varrho$ and $R(a_0, a_1), R(a_1, a_2), R(a_2, a_0)$, Definition 2 gives points $b_i, i > 0$, in \mathfrak{B} such that $R(b_0, b_1), R(b_1, b_2), R(b_2, b_3)$ and $b_3 \sim_{L, \varrho} b_0$. By the third and fourth lines in **(generation)**, we have $Z_1^w(b_3), Z^w(b_3)$, and so, by **(disjointness)**, $b_0 \neq b_3$. By the first formula in **(coordination)**, we must have $E_1(b_0, b_3)$.

The same argument as for b_0 above gives us an S -chain \bar{w} starting from b_3 and disjoint from \bar{v} in view of **(disjointness)** that defines uniquely an ω -word $w_{i'_1}w_{i'_2}\dots$ over Γ . The chain \bar{w} consists of consecutive segments $\bar{w}_{i'_1}, \bar{w}_{i'_2}, \dots$ such that $\bar{w}_{i'_1}$ sits in the same E_1 -equivalence class as \bar{v}_{i_1} , $\bar{w}_{i'_2}$ sits in an E_2 -equivalence class, $\bar{w}_{i'_3}$ in an E_1 -equivalence class, etc.

By the second formula in **(coordination)**, we have $i_1 = i'_1$, and so $(v_{i_1}, w_{i'_1}) \in \mathcal{P}$; by the third one, the ends of \bar{v}_{i_1} and $\bar{w}_{i'_1}$ are E_2 -related, from which $i_2 = i'_2$ and $(v_{i_2}, w_{i'_2}) \in \mathcal{P}$. The ends of \bar{v}_{i_2} and $\bar{w}_{i'_2}$ are E_1 -related, so $i_3 = i'_3$ and $(v_{i_3}, w_{i'_3}) \in \mathcal{P}$, etc.

Finally, let \bar{v} be $S(b_0, c_0), S(c_0, c_1), \dots$ and let \bar{w} be $S(b_3, d_0), S(d_0, d_1), \dots$. Since these S -chains are unique and $b_3 \sim_{L, \varrho} b_0$, we must have $c_i \sim_{L, \varrho} d_i$, for all $i < \omega$. It follows that the Γ -symbols carried by each pair c_i and d_i must coincide, which gives a solution to \mathcal{P} . \dashv

Since the ωPCP is undecidable, as an immediate consequence of Theorem 4 and Lemma 3 we obtain our main result:

Theorem 5. The $\text{FO}^2\text{E-IEP}$ and $\text{FO}^2\text{E-EDEP}$ are both undecidable.

4. Undecidability of the IEP for Guarded and Description Logics

It is not hard to tweak the construction in the previous section to show that the $\text{GF}^2\text{E}_c\text{-IEP}$ is also undecidable. Observe that the formula ψ defined above can be equivalently represented as a formula in GF^2E_c and that the $\text{FO}^2\text{E}(\varrho)$ -bisimulation between \mathfrak{A} and \mathfrak{B} is actually a $\text{GF}^2\text{E}_c(\varrho)$ -bisimulation (in this case, it is actually enough to have one copy of the R -cycle with an S -chain in \mathfrak{A}). The only unguarded conjunct of φ is $\forall x, y ((X_0(x) \wedge X_0(y)) \rightarrow x = y)$, which is needed to generate an R -cycle with three points. The same result can be achieved using a guarded φ' with two individual constants c_1 and c_2 as follows:

$$\varphi'(x) = R(x, c_1) \wedge R(c_1, c_2) \wedge R(c_2, x) \wedge \exists y [S(x, y) \wedge \bigwedge_{A_i \in \Gamma} (A_i(y) \rightarrow A_i(y))].$$

We then obtain Theorem 4 with φ' and GF^2E_c in place of φ and FO^2E , respectively, and, as an immediate consequence, the following:

Theorem 6. *The $\text{GF}^2\text{E}_c\text{-IEP}$ and $\text{GF}^2\text{E}_c\text{-EDEP}$ are both undecidable.*

These undecidability results can be interpreted as results about description logics with Boolean operators on roles that correspond to the two-variable fragments [36, 37]. The crucial operators required are role intersection, negation, and the identity role. Denote by $\mathcal{ALC}^{\cap, \neg, \text{id}}\text{E}$ the extension of the basic description logic \mathcal{ALC} with the operators \cap and \neg on roles, the identity role id interpreted as $\{(a, a) \mid a \in \text{dom}(\mathfrak{A})\}$ in any structure \mathfrak{A} , and two equivalence relations. The *interpolant existence problem* for $\mathcal{ALC}^{\cap, \neg, \text{id}}\text{E}$ (or $\mathcal{ALC}^{\cap, \neg, \text{id}}\text{E-IEP}$, for short) is the problem to decide, given $\mathcal{ALC}^{\cap, \neg, \text{id}}\text{E}$ -concepts C_1 and C_2 , whether there exists an $\mathcal{ALC}^{\cap, \neg, \text{id}}\text{E}$ -concept C built from the shared concept and role names from C_1 and C_2 such that $C_1 \sqsubseteq C$ and $C \sqsubseteq C_2$ are valid concept inclusions. We then obtain the following:

Theorem 7. *The $\mathcal{ALC}^{\cap, \neg, \text{id}}\text{E-IEP}$ and $\mathcal{ALC}^{\cap, \neg, \text{id}}\text{E-EDEP}$ are both undecidable.*

To obtain a DL version of the undecidability for the $\text{GF}^2\text{E}_c\text{-IEP}$, we can add nominals and the universal role to $\mathcal{ALC}^{\cap, \neg, \text{id}}\text{E}$ and, to reflect guarded quantification, admit only Boolean combinations of roles that contain at least one positive occurrence of a role name that is different from the universal role.

5. Open Problems

This paper presents first examples of decidable fragments of first-order logic, for which the interpolant existence problem and the explicit definition existence problem are undecidable. Numerous questions remain open, including the following:

- Is the IEP decidable for $\text{FO}^2\text{1E}$, that is, FO^2 with one equivalence relation? Note that the decidability and coNEXPTime -completeness proofs for $\text{FO}^2\text{1E}$ are significantly less involved than those for $\text{FO}^2\text{2E}$ [28]. What happens if we extend FO^2 with a single transitive (rather than equivalence) relation?
- Is the $\text{FO}^2\text{2E-IEP}$ still undecidable if one drops equality from FO^2 ? Note that, for FO^2 without equivalence relations and without equality, the IEP is decidable and the known complexity bounds are the same as for the case with equality [26].
- Is the IEP decidable for \mathcal{ALCQIO} ? What about C^2 , that is, FO^2 with counting quantifiers $\exists^{\geq k} x$, $k \in \mathbb{N}$? A possible hint that the IEP for C^2 might be undecidable is given by a recent result of [38] according to which the following related separation problem is undecidable: given two mutually exclusive C^2 -formulas φ and ψ , decide whether there exists an FO^2 -formula χ , called a *separator* for φ and ψ , such that $\varphi \models \chi$ and $\chi \models \neg\psi$.

References

- [1] K. L. McMillan, Interpolation and SAT-based model checking, in: Proceedings of the 15th International Conference on Computer Aided Verification, CAV 2003, Springer, 2003, pp. 1–13. URL: https://doi.org/10.1007/978-3-540-45069-6_1. doi:10.1007/978-3-540-45069-6_1.
- [2] R. Diaconescu, Logical support for modularisation, Logical environments 83 (1993) 130.
- [3] A. Cimatti, A. Griggio, R. Sebastiani, Interpolant generation for UTVPI, in: Proceedings of the 22nd International Conference on Automated Deduction, CADE 2009, Springer, 2009, pp. 167–182. URL: https://doi.org/10.1007/978-3-642-02959-2_15. doi:10.1007/978-3-642-02959-2_15.
- [4] D. Calvanese, S. Ghilardi, A. Gianola, M. Montali, A. Rivkin, Combined covers and Beth definability, in: Automated Reasoning - 10th International Joint Conference, IJCAR 2020, Paris, France, July 1-4, 2020, Proceedings, Part I, 2020, pp. 181–200. URL: https://doi.org/10.1007/978-3-030-51074-9_11. doi:10.1007/978-3-030-51074-9_11.
- [5] D. Calvanese, S. Ghilardi, A. Gianola, M. Montali, A. Rivkin, Combination of uniform interpolants via Beth definability, J. Autom. Reason. 66 (2022) 409–435. URL: <https://doi.org/10.1007/s10817-022-09627-1>. doi:10.1007/s10817-022-09627-1.
- [6] D. Toman, G. E. Weddell, Fundamentals of Physical Design and Query Compilation, Synthesis Lectures on Data Management, Morgan & Claypool Publishers, 2011.
- [7] M. Benedikt, J. Leblay, B. ten Cate, E. Tsamoura, Generating Plans from Proofs: The Interpolation-based Approach to Query Reformulation, Synthesis Lectures on Data Management, Morgan & Claypool Publishers, 2016.
- [8] B. ten Cate, E. Franconi, I. Seylan, Beth definability in expressive description logics, J. Artif. Intell. Res. 48 (2013) 347–414.
- [9] M. Benedikt, B. ten Cate, M. Vanden Boom, Effective interpolation and preservation in guarded logics, ACM Trans. Comput. Log. 17 (2016) 8. URL: <https://doi.org/10.1145/2814570>. doi:10.1145/2814570.
- [10] B. ten Cate, W. Conradie, M. Marx, Y. Venema, Definitorially complete description logics, in: Proc. of KR, 2006, pp. 79–89.
- [11] A. Artale, A. Mazzullo, A. Ozaki, F. Wolter, On free description logics with definite descriptions, in: Proceedings of the 18th International Conference on Principles of Knowledge Representation and Reasoning, KR 2021, 2021, pp. 63–73. URL: <https://doi.org/10.24963/kr.2021/7>. doi:10.24963/kr.2021/7.
- [12] D. Geleta, T. R. Payne, V. A. M. Tamma, An investigation of definability in ontology alignment, in: E. Blomqvist, P. Ciancarini, F. Poggi, F. Vitali (Eds.), Knowledge Engineering and Knowledge Management - 20th International Conference, EKAW 2016, Bologna, Italy, November 19-23, 2016, Proceedings, volume 10024 of *Lecture Notes in Computer Science*, 2016, pp. 255–271. URL: https://doi.org/10.1007/978-3-319-49004-5_17. doi:10.1007/978-3-319-49004-5_17.
- [13] B. Konev, C. Lutz, D. Walther, F. Wolter, Formal properties of modularisation, in: H. Stuckenschmidt, C. Parent, S. Spaccapietra (Eds.), Modular Ontologies: Concepts, Theories and Techniques for Knowledge Modularization, volume 5445 of *Lecture Notes in Computer Science*, Springer, 2009, pp. 25–66. URL: https://doi.org/10.1007/978-3-642-01907-4_3. doi:10.1007/978-3-642-01907-4_3.
- [14] E. Botoeva, B. Konev, C. Lutz, V. Ryzhikov, F. Wolter, M. Zakharyashev, Inseparability and conservative extensions of description logic ontologies: A survey, in: J. Z. Pan, D. Calvanese, T. Eiter, I. Horrocks, M. Kifer, F. Lin, Y. Zhao (Eds.), Reasoning Web: Logical Foundation of Knowledge Graph Construction and Query Answering - 12th International Summer School 2016, Aberdeen, UK, September 5-9, 2016, Tutorial Lectures, volume 9885 of *Lecture Notes in Computer Science*, Springer, 2016, pp. 27–89. URL: https://doi.org/10.1007/978-3-319-49493-7_2. doi:10.1007/978-3-319-49493-7_2.
- [15] I. Seylan, E. Franconi, J. de Bruijn, Effective query rewriting with ontologies over DBoxes, in: Proceedings of the 21st International Joint Conference on Artificial Intelligence, IJCAI 2009, 2009, pp. 923–925. URL: <http://ijcai.org/Proceedings/09/Papers/157.pdf>.

- [16] D. Toman, G. E. Weddell, First order rewritability for ontology mediated querying in Horn-DLFD, in: Proceedings of the 33rd International Workshop on Description Logics (DL 2020) co-located with the 17th International Conference on Principles of Knowledge Representation and Reasoning (KR 2020), Online Event [Rhodes, Greece], September 12th to 14th, 2020, 2020.
- [17] E. Franconi, V. Kerhet, N. Ngo, Exact query reformulation over databases with first-order and description logics ontologies, *J. Artif. Intell. Res.* 48 (2013) 885–922. URL: <https://doi.org/10.1613/jair.4058>. doi:10.1613/jair.4058.
- [18] E. Franconi, V. Kerhet, Effective query answering with ontologies and DBoxes, in: Description Logic, Theory Combination, and All That, Springer, 2019, pp. 301–328.
- [19] D. Toman, G. E. Weddell, FO Rewritability for OMQ using Beth Definability and Interpolation, in: Proceedings of the 34th International Workshop on Description Logics, DL 2021, CEUR-WS.org, 2021. URL: <http://ceur-ws.org/Vol-2954/paper-29.pdf>.
- [20] S. D. Comer, Classes without the amalgamation property., *Pacific J. Math.* 28 (1969) 309–318.
- [21] D. Pigozzi, Amalgamation, congruence-extension, and interpolation properties in algebras, *Algebra Univers.* (1971) 269–349.
- [22] M. Marx, C. Areces, Failure of interpolation in combined modal logics, *Notre Dame J. Formal Log.* 39 (1998) 253–273.
- [23] A. Artale, J. C. Jung, A. Mazzullo, A. Ozaki, F. Wolter, Living without Beth and Craig: Definitions and interpolants in description and modal logics with nominals and role inclusions, *ACM Trans. Comput. Log.* 24 (2023) 34:1–34:51. URL: <https://doi.org/10.1145/3597301>. doi:10.1145/3597301.
- [24] M. Fortin, B. Konev, F. Wolter, Interpolants and explicit definitions in extensions of the description logic \mathcal{EL} , in: Proceedings of the 19th International Conference on Principles of Knowledge Representation and Reasoning, KR 2022, 2022.
- [25] J. C. Jung, F. Wolter, Living without Beth and Craig: Definitions and interpolants in the guarded and two-variable fragments, in: Proceedings of the 36th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2021, IEEE, 2021, pp. 1–14. URL: <https://doi.org/10.1109/LICS52264.2021.9470585>. doi:10.1109/LICS52264.2021.9470585.
- [26] A. Kurucz, F. Wolter, M. Zakharyashev, Definitions and (uniform) interpolants in first-order modal logic, in: P. Marquis, T. C. Son, G. Kern-Isberner (Eds.), Proceedings of the 20th International Conference on Principles of Knowledge Representation and Reasoning, KR 2023, Rhodes, Greece, September 2-8, 2023, 2023, pp. 417–428. URL: <https://doi.org/10.24963/kr.2023/41>. doi:10.24963/kr.2023/41.
- [27] E. Kieronski, J. Michaliszyn, I. Pratt-Hartmann, L. Tendera, Two-variable first-order logic with equivalence closure, *SIAM J. Comput.* 43 (2014) 1012–1063. URL: <https://doi.org/10.1137/120900095>. doi:10.1137/120900095.
- [28] I. Pratt-Hartmann, Fragments of First-Order Logic, Oxford Logic Guides, Oxford University Press, United Kingdom, 2023.
- [29] V. Goranko, M. Otto, Model theory of modal logic, in: Handbook of Modal Logic, Elsevier, 2007, pp. 249–329.
- [30] E. Grädel, M. Otto, The freedoms of (guarded) bisimulation, in: Johan van Benthem on Logic and Information Dynamics, Springer International Publishing, 2014, pp. 3–31.
- [31] C. Chang, H. J. Keisler, Model Theory, Elsevier, 1998.
- [32] A. Robinson, A result on consistency and its application to the theory of definition, *Indagationes Mathematicae* 18 (1956) 47–58.
- [33] D. M. Gabbay, L. Maksimova, Interpolation and Definability: Modal and Intuitionistic Logics, Oxford, England: Oxford University Press UK, 2005.
- [34] K. Ruohonen, Reversible machines and Post’s correspondence problem for biprefix morphisms, *J. Inf. Process. Cybern.* 21 (1985) 579–595.
- [35] F. Gire, Two decidability problems for infinite words, *Inf. Process. Lett.* 22 (1986) 135–140. URL: [https://doi.org/10.1016/0020-0190\(86\)90058-X](https://doi.org/10.1016/0020-0190(86)90058-X). doi:10.1016/0020-0190(86)90058-X.
- [36] C. Lutz, U. Sattler, F. Wolter, Description logics and the two-variable fragment, in: C. A. Goble, D. L. McGuinness, R. Möller, P. F. Patel-Schneider (Eds.), Working Notes of the 2001

International Description Logics Workshop (DL-2001), Stanford, CA, USA, August 1-3, 2001, volume 49 of *CEUR Workshop Proceedings*, CEUR-WS.org, 2001. URL: <https://ceur-ws.org/Vol-49/LutzSattlerWolter-66start.ps>.

- [37] C. Lutz, U. Sattler, F. Wolter, Modal logic and the two-variable fragment, in: L. Fribourg (Ed.), *Computer Science Logic, 15th International Workshop, CSL 2001. 10th Annual Conference of the EACSL*, Paris, France, September 10-13, 2001, Proceedings, volume 2142 of *Lecture Notes in Computer Science*, Springer, 2001, pp. 247–261. URL: https://doi.org/10.1007/3-540-44802-0_18. doi:10.1007/3-540-44802-0_18.
- [38] L. Kuijter, T. Tan, F. Wolter, M. Zakharyashev, Separating counting from non-counting in fragments of two-variable first-order logic (extended abstract), in: *Proceedings of the 37th International Workshop on Description Logics*, CEUR Workshop Proceedings, CEUR-WS.org, 2024.